

**ACADEMIC SCIENTIFIC
JOURNAL OF CHEMISTRY**

ISSN: 2224-5286 (Print)
ISSN: 2518-1491 (Online)

**№3
2025**

ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)



ACADEMIC SCIENTIFIC JOURNAL OF CHEMISTRY

3 (464)

July – September 2025

PUBLISHED SINCE JANUARY

1947 PUBLISHED 4 TIMES A YEAR

ALMATY, NAS RK

Editor in chief:

ZHURINOV Murat Zhurinovich, doctor of chemistry, professor, academician of NAS RK, President of NAS RK RPA, general director of JSC "D.V. Sokolsky Institute of fuel, catalysis and electrochemistry (Almaty, Kazakhstan) <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

Editorial board:

ADEKENOV Sergazy Mynzhasarovich (deputy editor-in-chief) doctor of chemical sciences, professor, academician of NAS RK, director of the International Scientific and Production Holding «Phytochemistry» (Karaganda, Kazakhstan) <https://www.scopus.com/authid/detail.uri?authorId=7006153118>, <https://www.webofscience.com/wos/author/record/48648658>

AGABEKOV Vladimir Enokovich (deputy editor-in-chief), doctor of chemistry, professor, academician of NAS of Belarus, honorary director of the Institute of Chemistry of new materials (Minsk, Belarus) <https://www.scopus.com/authid/detail.uri?authorId=7004624845>, <https://www.webofscience.com/wos/author/record/28920574>

STRNAD Miroslav, head of the laboratory of the Institute of Experimental Botany of the Czech Academy of Sciences, professor (Olomouc, Czech Republic) <https://www.scopus.com/authid/detail.uri?authorId=36789185000>, <https://www.webofscience.com/wos/author/record/18379>

BURKITBAYEV Mukhambetkali, doctor of chemistry, professor, academician of NAS RK, (Almaty, Kazakhstan) <https://www.scopus.com/authid/detail.uri?authorId=8513885600>, <https://www.webofscience.com/wos/author/record/691218>

HOHMANN Judith, head of the department of pharmacognosy, faculty of Pharmacy, University of Szeged, director of the interdisciplinary center for Life sciences (Szeged, Hungary) <https://www.scopus.com/authid/detail.uri?authorId=7004457196>, <https://www.webofscience.com/wos/author/record/15630788>

ROSS Samir, Ph.D, professor, school of Pharmacy, National Center for scientific research of Herbal Products, University of Mississippi (Oxford, USA) <https://www.scopus.com/authid/detail.uri?authorId=7401610128>, <https://www.webofscience.com/wos/author/record/47926269>

KHUTORYANSKY Vitaly, Ph.D, pharmacist, professor at the University of Reading (Reading, England) <https://www.scopus.com/authid/detail.uri?authorId=35606915700>, <https://www.webofscience.com/wos/author/record/221621>

TELTAYEV Bagdat Burkhanbayuly, doctor of technical sciences, professor, academician of NAS RK, Ministry of Industry and infrastructure development of the Republic of Kazakhstan (Almaty, Kazakhstan) <https://www.scopus.com/authid/detail.uri?authorId=6506225641>, <https://www.webofscience.com/wos/author/record/72161>

PHARUK Asana Dar, professor at Hamdard al-Majid college of Oriental medicine, faculty of Oriental medicine, Hamdard University (Karachi, Pakistan) <https://www.scopus.com/authid/detail.uri?authorId=55884056900>, <https://www.webofscience.com/wos/author/record/1796996>

FAZYLOV Serik Drakhmetovich, doctor of chemistry, professor, academician of NAS RK, deputy director of the Institute of Organic Synthesis and Coal Chemistry (Karaganda, Kazakhstan) <https://www.scopus.com/authid/detail.uri?authorId=6701472056>, <https://www.webofscience.com/wos/author/record/1541357>

ZHOROBEKOVA Sharipa Zhorobekovna, doctor of chemistry, professor, academician of NAS of Kyrgyzstan, Institute of Chemistry and chemical technology of NAS KR (Bishkek, Kyrgyzstan) <https://www.scopus.com/authid/detail.uri?authorId=6602652060>, <https://www.webofscience.com/wos/author/record/31723468>

KHALIKOV Jurabay Khalikovich, doctor of chemistry, professor, academician of the Academy of Sciences of Tajikistan, V.I. Nikitin Institute of Chemistry AS RT (Tajikistan) <https://www.scopus.com/authid/detail.uri?authorId=6603735641>, <https://www.webofscience.com/wos/author/record/9567106>

FARZALIEV Vagif Medzhid ogly, doctor of chemistry, professor, academician of NAS of Azerbaijan (Azerbaijan) <https://www.scopus.com/authid/detail.uri?authorId=6601962486>, <https://www.webofscience.com/wos/author/record/21617033>

GARELIK Hemda, PhD in chemistry, president of the department of Chemistry and Environment of the International Union of Pure and Applied Chemistry (London, England) <https://www.scopus.com/authid/detail.uri?authorId=56010090400>, <https://www.webofscience.com/wos/author/record/29866743>

ACADEMIC SCIENTIFIC JOURNAL OF CHEMISTRY

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Owner: «Central Asian Academic Research Center» LLP (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Social Development of the Republic of Kazakhstan № **KZ23VPY00121156**, issued 05.06.2025

Thematic scope: *organic chemistry, inorganic chemistry, catalysis, electrochemistry and corrosion, pharmaceutical chemistry and technology.*

Periodicity: 4 times a year.

<http://chemistry-technology.kz/index.php/en/arhiv>

© «Central Asian Academic Research Center» LLP, 2025

Editorial address: JSC «D.V. Sokolsky institute of fuel, catalysis and electrochemistry», 142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22, e-mail: orgcat@nursat.kz

Бас редактор:

ЖҰРЫНОВ Мұрат Жұрыңұлы, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, РБҚ ҚР ҰҒА президенті, АҚ «Д.В. Сокольский атындағы Отын, катализ және электрохимия институтының» бас директоры (Алматы, Қазақстан) <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

Редакция алқасы:

ӘДЕКЕНОВ Серғазы Мыңжасарұлы (бас редактордың орынбасары), химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, «Фитохимия» Халықаралық ғылыми-өндірістік холдингінің директоры (Қарағанды, Қазақстан) <https://www.scopus.com/authid/detail.uri?authorId=7006153118>, <https://www.webofscience.com/wos/author/record/48648658>

АГАБЕКОВ Владимир Енокович (бас редактордың орынбасары), химия ғылымдарының докторы, профессор, Беларусь ҰҒА академигі, Жаңа материалдар химиясы институтының құрметті директоры (Минск, Беларусь) <https://www.scopus.com/authid/detail.uri?authorId=7004624845>, <https://www.webofscience.com/wos/author/record/28920574>

СТРНАД Мирослав, профессор, Чехия ғылым академиясының Эксперименттік ботаника институтының зертхана меңгерушісі (Оломоуц, Чехия) <https://www.scopus.com/authid/detail.uri?authorId=36789185000>, <https://www.webofscience.com/wos/author/record/18379>

БҮРКІТБАЕВ Мұхамбетқали, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, (Алматы, Қазақстан) <https://www.scopus.com/authid/detail.uri?authorId=8513885600>, <https://www.webofscience.com/wos/author/record/691218>

ХОХМАНН Джулит, Сегед университетінің Фармацевтика факультетінің Фармакогнозия кафедрасының меңгерушісі, Жаратылыстану ғылымдарының пәнаралық орталығының директоры (Сегед, Венгрия) <https://www.scopus.com/authid/detail.uri?authorId=7004457196>, <https://www.webofscience.com/wos/author/record/15630788>

РОСС Самир, PhD, Миссисипи университетінің Өсімдік өнімдерін ғылыми зерттеу ұлттық орталығы, Фармация мектебінің профессоры (Оксфорд, АҚШ) <https://www.scopus.com/authid/detail.uri?authorId=7401610128>, <https://www.webofscience.com/wos/author/record/47926269>

ХУТОРЯНСКИЙ Виталий, философия докторы (PhD, фармацевт), Реддинг университетінің профессоры (Реддинг, Англия) <https://www.scopus.com/authid/detail.uri?authorId=35606915700>, <https://www.webofscience.com/wos/author/record/221621>

ТЕЛТАЕВ Бағдат Бұрханбайұлы, техника ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Қазақстан Республикасы Индустрия және инфрақұрылымдық даму министрлігі (Алматы, Қазақстан) <https://www.scopus.com/authid/detail.uri?authorId=6506225641>, <https://www.webofscience.com/wos/author/record/72161>

ФАРУК Асана Дар, Хамдар аль-Маджида Шығыс медицина колледжінің профессоры, Хамдард университетінің Шығыс медицина факультеті (Карачи, Пәкістан) <https://www.scopus.com/authid/detail.uri?authorId=55884056900>, <https://www.webofscience.com/wos/author/record/1796996>

ФАЗЫЛОВ Серік Драхметұлы, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Органикалық синтез және көмір химиясы институты директорының ғылыми жұмыстар жөніндегі орынбасары (Қарағанды, Қазақстан) <https://www.scopus.com/authid/detail.uri?authorId=6701472056>, <https://www.webofscience.com/wos/author/record/1541357>

ЖОРОБЕКОВА Шарипа Жоробекқызы, химия ғылымдарының докторы, профессор, Кыргызстан ҰҒА академигі, ҚР ҰҒА Химия және химиялық технология институты (Бішкек, Кыргызстан) <https://www.scopus.com/authid/detail.uri?authorId=6602652060>, <https://www.webofscience.com/wos/author/record/31723468>

ХАЛИКОВ Джурабай Халикович, химия ғылымдарының докторы, профессор, Тәжікстан ҒА академигі, В.И. Никитин атындағы Химия институты (Душанбе, Тәжікстан) <https://www.scopus.com/authid/detail.uri?authorId=6603735641>, <https://www.webofscience.com/wos/author/record/9567106>

ФАРЗАЛИЕВ Вагиф Меджидоглы, химия ғылымдарының докторы, профессор, АҰҒА академигі (Баку, Әзірбайжан) <https://www.scopus.com/authid/detail.uri?authorId=6601962486>, <https://www.webofscience.com/wos/author/record/21617033>

ГАРЕЛИК Хемда, философия докторы (PhD, химия), Халықаралық таза және қолданбалы химия одағының Химия және қоршаған орта бөлімінің президенті (Лондон, Англия) <https://www.scopus.com/authid/detail.uri?authorId=56010090400>, <https://www.webofscience.com/wos/author/record/29866743>

«ACADEMIC SCIENTIFIC JOURNAL OF CHEMISTRY»

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Меншіктенуші: «Орталық Азия академиялық ғылыми орталығы» ЖШС (Алматы қ.).

Қазақстан Республикасының Ақпарат және қоғамдық даму министрлігінің Ақпарат комитетінде 05.06.2025 ж. берілген № KZ23VPY00121156 мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *органикалық химия, бейорганикалық химия, катализ, электрохимия және коррозия, фармацевтикалық химия және технологиялар.*

Мерзімділігі: жылына 4 рет.

<http://chemistry-technology.kz/index.php/en/arhiv>

© «Орталық Азия академиялық ғылыми орталығы» ЖШС, 2025

Редакцияның мекенжайы: 050100, Алматы қ., Қонаев к-сі, 142, «Д.В. Сокольский атындағы отын, катализ және электрохимия институты» АҚ, каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Главный редактор:

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик НАН РК, президент РОО Национальной академии наук Республики Казахстан, генеральный директор АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского» (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>, <https://www.webofscience.com/wos/author/record/2017489>

Редакционная коллегия:

АДЕКЕНОВ Сергазы Мынжасарович (заместитель главного редактора), доктор химических наук, профессор, академик НАН РК, директор Международного научно-производственного холдинга «Фитохимия» (Караганда, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=7006153118>, <https://www.webofscience.com/wos/author/record/48648658>

АГАБЕКОВ Владимир Енокович (заместитель главного редактора), доктор химических наук, профессор, академик НАН Беларуси, почетный директор Института химии новых материалов (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>, <https://www.webofscience.com/wos/author/record/28920574>

СТРНАД Мiroслав, профессор, заведующий лабораторией института Экспериментальной ботаники Чешской академии наук (Оломоуц, Чехия), <https://www.scopus.com/authid/detail.uri?authorId=36789185000>, <https://www.webofscience.com/wos/author/record/18379>

БУРКИТБАЕВ Мухамбеткали, доктор химических наук, профессор, академик НАН РК, (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=8513885600>, <https://www.webofscience.com/wos/author/record/691218>

ХОХМАНН Джудит, заведующий кафедрой Фармакогнозии Фармацевтического факультета Университета Сегеда, директор Междисциплинарного центра естественных наук (Сегед, Венгрия), <https://www.scopus.com/authid/detail.uri?authorId=7004457196>, <https://www.webofscience.com/wos/author/record/15630788>

РОСС Самир, PhD, профессор Школы Фармации национального центра научных исследований растительных продуктов Университета Миссисипи (Оксфорд, США), <https://www.scopus.com/authid/detail.uri?authorId=7401610128>, <https://www.webofscience.com/wos/author/record/47926269>

ХУТОРЯНСКИЙ Виталий, доктор философии (Ph.D, фармацевт), профессор Университета Рединга (Рединг, Англия), <https://www.scopus.com/authid/detail.uri?authorId=35606915700>, <https://www.webofscience.com/wos/author/record/221621>

ТЕЛЫТАЕВ Багдат Бурханбайулы, доктор технических наук, профессор, академик НАН РК, Министерство Индустрии и инфраструктурного развития Республики Казахстан (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6506225641>, <https://www.webofscience.com/wos/author/record/72161>

ФАРУК Ахсана Дар, профессор колледжа Восточной медицины Хамдарда аль-Маджида, факультет Восточной медицины университета Хамдарда (Карачи, Пакистан), <https://www.scopus.com/authid/detail.uri?authorId=55884056900>, <https://www.webofscience.com/wos/author/record/1796996>

ФАЗЫЛОВ Серик Драгметович, доктор химических наук, профессор, академик НАН РК, заместитель директора по научной работе Института органического синтеза и углехимии (Караганда, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6701472056>, <https://www.webofscience.com/wos/author/record/1541357>

ЖОРОБЕКОВА Шарипа Жоробековна, доктор химических наук, профессор, академик НАН Кыргызстана, Институт химии и химической технологии НАН КР (Бишкек, Кыргызстан), <https://www.scopus.com/authid/detail.uri?authorId=6602652060>, <https://www.webofscience.com/wos/author/record/31723468>

ХАЛИКОВ Джурабай Халикович, доктор химических наук, профессор, академик АН Таджикистана, Институт химии имени В.И. Никитина АН РТ (Душанбе, Таджикистан), <https://www.scopus.com/authid/detail.uri?authorId=6603735641>, <https://www.webofscience.com/wos/author/record/9567106>

ФАРЗАЛИЕВ Вагиф Меджид оглы, доктор химических наук, профессор, академик НАНА (Баку, Азербайджан), <https://www.scopus.com/authid/detail.uri?authorId=6601962486>, <https://www.webofscience.com/wos/author/record/21617033>

ГАРЕЛИК Хемда, доктор философии (Ph.D, химия), президент Отдела химии и окружающей среды Международного союза чистой и прикладной химии (Лондон, Англия), <https://www.scopus.com/authid/detail.uri?authorId=56010090400>, <https://www.webofscience.com/wos/author/record/29866743>

«ACADEMIC SCIENTIFIC JOURNAL OF CHEMISTRY».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Собственник: Республиканское общественное объединение ТОО «Центрально-азиатский академический научный центр» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и общественного развития Республики Казахстан № KZ23VPY00121156, выданное 05.06.2025 г.

Тематическая направленность: *органическая химия, неорганическая химия, катализ, электрохимия и коррозия, фармацевтическая химия и технологии.*

Периодичность: 4 раз в год.

<http://chemistry-technology.kz/index.php/en/arhiv>

© ТОО «Центрально-азиатский академический научный центр», 2025

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142, АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского», каб. 310, тел. 291-62-80, факс 291-57-22, e-mail: orgcat@nursat.kz

CONTENTS

A.D. Alakhunova, L.E. Agibayeva, R.A. Mangazbayeva Obtaining of methylcellulose-based hydrogels using radiation treatment method.....	11
A.P. Auyeshov, K.T. Arynov, Ch.Z. Yeskibayeva Resource and economic efficiency of serpentinite waste utilization for the production of inorganic magnesium compounds.....	29
S. Duzelbayeva, B. Imangaliyeva, A. Aldiyarova, N. Sovet, B. Bakhtiyarov Production of biopolymer from starch as an alternative to artificial polymer and study of its biodegradable properties.....	41
G.D. Jetpisbayeva, B.K. Massalimova, V.A. Sadykov, A. Darmenbayeva, G.B. Aubakirova The influence of production methods on the selectivity and stability of Co-containing catalysts for Fischer-Tropsch synthesis.....	64
B.K. Kenzhaliyev, A.K. Koizhanova, T.Yu. Surkova, Z.D. Dosymbayeva, D.M. Yessimova Investigation of the modes and parameters of gold leaching from man-made raw materials.....	75
M.K. Kurmanaliev, Zh.E. Shaikhova, L.M. Kalimoldina, S.O. Abilkasova Synthesis and sorption properties of new selective sorbents based on crown ethers...	92
E. Kairatuly, E.K. Assembayeva, A.Zh Bozhbanov, D.E. Nurmukhanbetova, E.Zh. Gabdullina Evaluation of physicochemical indicators of combined fermented milk products....	102
A.B. Kuandykova, B.Zh. Dzhiembraev, N.I. Akylbekov, A.B. Dobrynin Molecular and crystal structures of 4-dimethoxyphosphoryltetrahydropyran (thiopyran)-4-ol.....	115
R.M. Kudaibergenova, S.A. Orynbayev, E.A. Baibazarova, K.B. Bulekbayeva, G.A. Seitbekova Technological pathways for sustainable wastewater treatment.....	127
G.M. Madybekova, A.B. Issayeva, B.Zh. Mutaliyeva, S.S. Bitursyn Physicochemical properties of microcapsules based on natural polymers containing probiotic microorganisms.....	140
M.M. Mataev, Z.B. Sarsenbayeva, M.A. Nurbekova, M.R. Abdraimova, K.Zh. Seitbekova Synthesis and morphological analysis of $\text{Fe}_{0.84}\text{Mn}_{1.12}\text{O}_3\text{-In}_{0.12}\text{Fe}_{1.88}\text{O}_3$ composite.....	155

Z. Muldakhmetov, S. Fazylov, O. Nurkenov, Zh. Akhmetkarimova, O. Seilkhanov Synthesis and properties of new naphthyl-containing thiosemicarbazides and thioureas.....	166
U. Nazarbek, P. Abdurazova, G. Kambarova, Y. Raiymbekov Machine learning-based prediction of temperature-driven solubility changes in aqueous salt solutions.....	184
R.S. Orazbekova, S.A. Tungatarova, A.E. Tolembek, A.O. Aidarova, M.K. Yerkibaeva Catalytic processing of renewable raw materials into hydrogen-containing fuel mixtures.....	194
S.K. Rakhimova, R.I. Jalmakhanbetova, G.K. Mukusheva, A.A. Asylbekova, Zh. Zh. Zhumagaliyeva Spectroscopic analysis of methanol extract of <i>Ziziphora bungeana</i> Juz. and study of its antibacterial activity.....	207
R.K. Rakhmetullaeva, B. Khavilkhairat, N.B. Sarova, G.O. Rvaidarova, A.N. Nurlybayeva Copolymers based on acrylic acid for water purification from heavy metal ions.....	219
A.N. Sabitova, Zh.S. Kassymova, R.E. Mukiyanova, B.B. Bayahmetova, N.N. Nurgaliyev Investigation of the effectiveness of metallurgical slags in fertilizer production.....	233
E.T. Talgatov, D.A. Bibatyrova, A.A. Naizabaev, S.A. Kuttybayeva, A.Z. Abilmagzhanov Selective hydrogenation of phenylacetylene over polymer-modified Pd catalysts immobilized on inorganic supports.....	243
S. Tyanakh, T.O. Khamitova, A.P. Nauanova, D.M. Kereevna, A.S. Darmenbayeva Study of the properties of humic acids synthesized from brown coal of the Kuznetsk and the Kumuskuduk deposits.....	255
A.A. Turgunbaeva, G.N. Gemejiyeva, N.A. Sultanova Investigation of the chemical composition of the chloroform extract of <i>Rheum Tataricum</i> L. f. By gas-chromatography.....	275

МАЗМҰНЫ

Ә.Д. Алахунова, Л.Ә. Агибаева, Р.А. Мангазбаева Метилцеллюлоза негізінде гидрогельдерді радиациялық өңдеу әдісімен алу.....	11
А.П. Ауешов, К.Т. Арынов, Ч.З. Ескибаева Магнийдің бейорганикалық қосылыстарын алу мақсатында серпентинитті қайта өңдеудің ресурстық және экономикалық тиімділігі.....	29
С. Дузелбаева, Б. Иманғалиева, А. Алдиярова, Н. Совет, Б. Бақтияров Жасанды полимерге балама ретінде крахмалдан биополимер алу және оның биоыдырау қасиеттерін зерттеу.....	41
Г.Д. Джетписбаева, Б.К. Масалимова, В.А. Садыков, А. Дарменбаева, Г.Б. Аубакирова Фишер-Тропш синтезінің Со-құрамды катализаторларының талғамдылығы мен тұрақтылығына дайындау әдістерінің әсері.....	64
Б.К. Кенжалиев, А.К. Койжанова, Т.Ю. Суркова, З.Д. Досымбаева, Д.М. Есимова Техногендік шикізаттан алтынды сілтілеудің режимдері мен параметрлерін зерттеу.....	75
М.Қ. Құрманалиев, Ж.Е. Шаихова, Л.М. Калимолдина, С.О. Әбілқасова Краун-эфирлер негізіндегі жаңа талғамды сорбенттердің синтезі мен сорбциялық қасиеттері.....	92
Е. Қайратұлы, Э. К. Асембаева, А.Ж. Божбанов, Д.Е. Нурмуханбетова, Е.Ж. Габдуллина Құрамдастырылған сүтқышқылды өнімдердің физика-химиялық көрсеткіштерін бағалау.....	102
А.Б. Қуандықова, Б.Ж. Джиембаев, Н.И. Акылбеков, А.Б. Добрынин 4-Диметоксифосфорилтетрагидропиран(тиопиран)-4-олдардың молекулалық және кристалдық құрылымдары.....	115
Р.М. Құдайбергенова, С.А. Орынбаев, Е.А. Байбазарова, Қ.Б. Бөлекбаева, Г.А. Сейтбекова Ағынды суларды тұрақты тазартудың технологиялық жолдары.....	127
Г.М. Мадыбекова, А.Б. Исаева, Б.Ж. Муталиева, С.С. Битурсын Табиғи полимерлер негізіндегі, пробиотикалық микроорганизмдер қамтылған микрокапсулалардың физика-химиялық қасиеттері.....	140

М.М. Матаев, З.Б. Сарсенбаева, М.А. Нурбекова, М.Р. Абдраимова, К.Ж. Сейтбекова $\text{Fe}_{0.84}\text{Mn}_{1.12}\text{O}_3\text{-IN}_{0.12}\text{Fe}_{1.88}\text{O}_3$ композитінің синтезі және морфологиялық талдауы.....	155
З. Молдахметов, С. Фазылов, О. Нүркенов, Ж. Ахметкәрімова, О. Сейілханов Жаңа нафтилді тиосемикарбазидтер мен тиомочевиналардың синтезі мен қасиеттері.....	166
У. Назарбек, П. Абдуразова, Ғ. Қамбарова, Е. Райымбеков Сулы ерітінділердегі тұздардың ерігіштігінің температуралық өзгерістерін машиналық оқыту әдістерімен болжау.....	184
Р.С. Оразбекова, С.А. Тунгатарова, А.Е. Төлембек, А.О. Айдарова, М.Қ. Еркібаева Жаңартылатын шикізатты құрамында сутегі бар отын қоспаларына дейін каталитикалық өңдеу.....	194
С.Қ. Рахимова, Р.И. Джалмаханбетова, Г.К. Мукушева, А.А. Асылбекова, Ж.Ж. Жумағалиева <i>Ziziphora Bungeana</i> Juz. метанолды сығындысын спектроскопиялық талдау және оның бактерияға қарсы белсенділігін зерттеу.....	207
Р.Қ. Рахметуллаева, Б. Хавилхайрат, Н.Б. Сарова, Г.О. Рвайдарова, А.Н. Нурлыбаева Ауыр металл иондарынан су тазалауға арналған акрил қышқылы негізіндегі сополимерлер.....	219
А.Н. Сабитова, Ж.С. Касымова, Р.Е. Мукиянова, Б.Б. Баяхметова, Н.Н. Нургалиев Тыңайтқыштар өндірісіндегі металлургиялық шлактардың тиімділігін зерттеу.....	233
Э.Т. Талғатов, Д.А. Бибатырова, А.А. Найзабаев, Ш.Ә. Құттыбаева, А.З. Абильмагжанов Бейорганикалық тіректерде иммобилизацияланған полимермен модификацияланған PD катализаторлары бойынша фенилацетиленді селективті гидрогенизациялау.....	243
С. Тянах, Т.О. Хамитова, А.П. Науанова, Д.М. -Кереевна, А.С. Дарменбаева Кузнецк және Күмісқұдық қоңыр көмірінен синтезделіп алынатын гумин қышқылдарының қасиеттерін зерттеу.....	255
А.А. Тургунбаева, Н.Г. Гемеджиева, Н.А. Султанова <i>Rheum Tataricum</i> L. f. өсімдігінің хлороформ сығындысының химиялық құрамын газ хроматография әдісімен зерттеу.....	275

СОДЕРЖАНИЕ

A.D. Alakhunova, L.E. Agibayeva, R.A. Mangazbayeva Получение гидрогелей на основе метилцеллюлозы методом радиационной обработки.....	11
A.P. Auyeshov, K.T. Arynov, Ch.Z. Yeskibayeva Ресурсная и экономическая эффективность утилизации отходов серпентинита для производства неорганических соединений магния.....	29
S. Duzelbayeva, B. Imangaliyeva, A. Aldiyarova, N. Sovet, B. Bakhtiyarov Получение биополимера из крахмала как альтернатива искусственному полимеру и исследование его биоразлагаемых свойств.....	41
G.D. Jetpisbayeva, B.K. Massalimova, V.A. Sadykov, A. Darmenbayeva, G.B. Aubakirova Влияние методов получения на селективность и стабильность катализаторов, содержащих кобальт, для синтеза по Фишеру-Тропшу.....	64
B.K. Kenzhaliyev, A.K. Koizhanova, T.Yu. Surkova, Z.D. Dosymbayeva, D.M. Yessimova Исследование режимов и параметров выщелачивания золота из техногенного сырья.....	75
M.K. Kurmanaliev, Zh.E. Shaikhova, L.M. Kalimoldina, S.O. Abilkasova Синтез и сорбционные свойства новых селективных сорбентов на основе краун-эфиров.....	92
E. Kairatuly, E.K. Assembayeva, A.Zh. Bozhbanov, D.E. Nurmukhanbetova, E.Zh. Gabdullina Оценка физико-химических показателей комбинированных кисломолочных продуктов.....	102
A.B. Kuandykova, B.Zh. Dzhiembaev, N.I. Akylbekov, A.B. Dobrynin Молекулярные и кристаллические структуры 4-диметоксифосфорилтетрагидропиран(тиопиран)-4-ола.....	115
R.M. Kudaibergenova, S.A. Orynbayev, E.A. Baibazarova, K.B. Bulekbayeva, G.A. Seitbekova Технологические пути устойчивой очистки сточных вод.....	127
G.M. Madybekova, A.B. Issayeva, B.Zh. Mutaliyeva, S.S. Bitursyn Физико-химические свойства микрокапсул на основе природных полимеров, содержащих пробиотические микроорганизмы.....	140

M.M. Mataev, Z.B. Sarsenbayeva, M.A. Nurbekova, M.R. Abdraimova, K.Zh. Seitbekova Синтез и морфологический анализ композита Fe _{0.84} Mn _{1.12} O ₃ -In _{0.12} Fe _{1.88} O ₃	155
Z. Muldakhmetov, S. Fazylov, O. Nurkenov, Zh. Akhmetkarimova, O. Seilkhanov Синтез и свойства новых нафтилилсодержащих тиосемикарбазидов и тиомочевин.....	166
U. Nazarbek, P. Abdurazova, G. Kambarova, Y. Raiymbekov Прогнозирование изменений растворимости солей в воде в зависимости от температуры с использованием машинного обучения.....	184
R.S. Orazbekova, S.A. Tungatarova, A.E. Tolembek, A.O. Aidarova, M.K. Yerkibaeva Каталитическая переработка возобновляемого сырья в водородсодержащие топливные смеси.....	194
S.K. Rakhimova, R.I. Jalmakhanbetova, G.K. Mukusheva, A.A. Asylbekova, Zh. Zh. Zhumagaliyeva Спектроскопический анализ метанольного экстракта <i>Ziziphora bungeana</i> Juz. и исследование его антибактериальной активности.....	207
R.K. Rakhmetullaeva, B. Khavilkhairat, N.B. Sarova, G.O. Rvaidarova, A.N. Nurlybayeva Кополимеры на основе акриловой кислоты для очистки воды от ионов тяжёлых металлов.....	219
A.N. Sabitova, Zh.S. Kassymova, R.E. Mukiyanova, B.B. Bayahmetova, N.N. Nurgaliev Исследование эффективности металлургических шлаков при производстве удобрений.....	233
E.T. Talgatov, D.A. Bibatyrova, A.A. Naizabaev, S.A. Kuttybayeva, A.Z. Abilmagzhanov Селективное гидрирование фенилэтина на модифицированных полимерами PD-катализаторах, иммобилизованных на неорганических носителях.....	243
S. Tyanakh, T.O. Khamitova, A.P. Nauanova, D.M. Kereevna, A.S. Darmenbayeva Исследование свойств гуминовых кислот, синтезированных из бурого угля Кузнецкого и Кумускудукского месторождений.....	255
A.A. Turgunbaeva, G.N. Gemejiyeva, N.A. Sultanova Исследование химического состава хлороформного экстракта <i>Rheum</i> <i>Tataricum</i> L. f. методом газовой хроматографии.....	275

<https://doi.org/10.32014/2025.2518-1491.315>

IRSTI 52.35.01

UDC 622.33; 622.337.2

© S. Tyanakh^{1*}, T.O. Khamitova², A.P. Nauanova², D.M. Kereevna³,
A.S. Darmenbayeva⁴, 2025.

¹E.A. Buketov Karagandy University, Karaganda, Kazakhstan;

²NCJSC «S. Seifullin Kazakh Agro Technical Research University»,
Astana, Kazakhstan;

³Al-Farabi Kazakh National University, Almaty, Kazakhstan;

⁴M.Kh. Dulaty Taraz University, Taraz, Kazakhstan.

E-mail: saika_8989@mail.ru

STUDY OF THE PROPERTIES OF HUMIC ACIDS SYNTHESIZED FROM BROWN COAL OF THE KUZNETSK AND THE KUMUSKUDUK DEPOSITS

Tyanakh Sairagul — PhD, E.A. Buketov Karagandy University, Karaganda, Kazakhstan,

E-mail: saika_8989@mail.ru; <https://orcid.org/0000-0001-5343-4695>;

Khamitova Tolkyn Ondirisovna — PhD, acting associate professor, NCJSC «S.Seifullin Kazakh Agro Technical Research University», Astana, Kazakhstan,

E-mail: khamitova.t@inbox.ru; <https://orcid.org/0000-0002-4691-3732>;

Nauanova Ainash Pakhuashovna — Doctor of Biological Sciences, Professor, NCJSC «S.Seifullin Kazakh Agro Technical Research University», Astana, Kazakhstan,

E-mail: nauanova@mail.ru; <https://orcid.org/0000-0002-5250-1961>;

Dana Mykty-Kereevna Ibraimova — Candidate of Chemical Sciences (CSc), Senior Lecturer, Department of analytical, colloid chemistry and technology of rare elements, Faculty of Chemistry and Chemical technology, Al-Farabi Kazakh National University», Almaty, Kazakhstan,

E-mail: dana_kereevna@kaznu.kz <https://orcid.org/0000-0002-1761-9745>;

Darmenbayeva Akmaral Sabetbekkyzy — Associate professor at the Department of Chemistry and Chemical technology, M.Kh. Dulaty Taraz University, Taraz, Kazakhstan,

E-mail: maral88@mail.ru, ORCID: <https://orcid.org/0000-0003-2974-0398>.

Abstract. The article is devoted to the synthesis and comparative analysis of humic acids derived from brown coal from the Kuznetsk and Kumuskuduk deposits. Using advanced analytical techniques, including infrared (IR) spectroscopy, elemental analysis, and thermogravimetric analysis, we evaluated the physicochemical properties of humic acids from these two geological sources. The research revealed significant differences in chemical composition, functional groups, and thermal stability, attributed to the distinct geological formation conditions of each deposit. In particular, variations in carboxyl and hydroxyl group content were observed, impacting the sorption and complexation capabilities of the humic acids. This comparative study highlights the potential of humic acids from both deposits for applications in environmental remediation, agriculture, and

wastewater treatment. The findings provide new insights into the influence of geological factors on the structural characteristics and reactivity of humic acids, thus expanding the possibilities for their practical use and further technological development. This allows us to assess their potential application in various industries, such as agrochemistry, ecology and medicine. Thus, the research results obtained are a significant contribution to the field of coal chemistry and ecology, and can also serve as the basis for further research and development in the field of the use of humic acids as environmentally friendly additives and sorbents.

Keywords: brown coal, humic acids, Kumuskuduk deposit, Kuznetsk deposit, functional groups

Financing. *This research was conducted as part of the targeted funding program for the project IRN BR24992961, titled “Development of new technologies using biological systems for processing coal waste into organo-mineral fertilizer to increase soil fertility and crop productivity”.*

© С. Тянах^{1*}, Т.О. Хамитова², А.П. Науанова², Д.М. -Кереевна³,
А.С. Дарменбаева⁴, 2025.

¹Е.А. Бөкетов атындағы Қарағанды университеті, Қарағанды, Қазақстан;

²Сейфуллин атындағы Қазақ агротехникалық зерттеу университеті,
Астана, Қазақстан;

³Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан;

⁴М.Х. Дулати атындағы Тараз университеті, Тараз, Қазақстан.

E-mail: saika_8989@mail.ru

КУЗНЕЦК ЖӘНЕ КҮМІСКҰДЫҚ ҚОҢЫР КӨМІРІНЕН СИНТЕЗДЕЛІП АЛЫНАТЫН ГУМИН ҚЫШҚЫЛДАРЫНЫҢ ҚАСИЕТТЕРІН ЗЕРТТЕУ

Тянах Сайрагул — PhD, академик Е.А. Букетов атындағы Қарағанды университеті, Қарағанды, Қазақстан,

E-mail: saika_8989@mail.ru, <https://orcid.org/0000-0001-5343-4695>;

Хамитова Толқын Ондирисовна — PhD, агрономия факультеті, С. Сейфуллин атындағы Қазақ агротехникалық университеті, Астана, Қазақстан,

E-mail: khamitova.t@inbox.ru, <https://orcid.org/0000-0002-4691-3732>;

Науанова Айнаш Пахуашовна — биология ғылымдарының докторы, профессор, «С. Сейфуллин атындағы Қазақ агротехникалық зерттеу университеті» КеАҚ, Астана, Қазақстан, E-mail: nauanova@mail.ru, <https://orcid.org/0000-0002-5250-1961>;

Дана Мықты-Кереевна Ибраимова — химия ғылымдарының кандидаты, Әл-Фараби атындағы Қазақ ұлттық университеті, Химия және химиялық технология факультеті, Аналитикалық, коллоидтық химия және сирек элементтер технологиясы кафедрасының аға оқытушысы, Алматы, Қазақстан,

E-mail: dana_kereevna@kaznu.kz, ORCID: <https://orcid.org/0000-0002-1761-9745>;

Дарменбаева Акмарал Сабетбекқызы — М.Х. Дулати атындағы Тараз университеті, Химия және химиялық технология кафедрасының доценті, Тараз, Қазақстан,

E-mail: maral88@mail.ru; <https://orcid.org/0000-0003-2974-0398>.

Аннотация. Бұл зерттеу Кузнецк кен орыны және Күміскұдық кен орындарының қоңыр көмірінен алынған гумин қышқылдарының синтезі мен салыстырмалы талдауына арналады. Жетілдірілген аналитикалық әдістерді, соның ішінде инфрақызыл спектроскопиялық талдау, элементтік талдау және термогравиметриялық талдау әдістерін пайдалана отырып, біз осы екі геологиялық көздерден гумин қышқылдарының физика-химиялық қасиеттерін бағаладық. Зерттеу нәтижесінде әрбір кен орнының геологиялық түзілу жағдайларына байланысты химиялық құрамы, функционалдық топтары және термиялық тұрақтылығы бойынша елеулі айырмашылықтар анықталды. Атап айтқанда, гумин қышқылдарының сорбциялық және комплекс түзу мүмкіндіктеріне әсер ететін карбоксил және гидроксил тобының құрамындағы өзгерістер байқалды. Бұл салыстырмалы зерттеу қоршаған ортаны қалпына келтіруде, ауыл шаруашылығында және ағынды суларды тазартуда қолдану үшін екі кен орнының гумин қышқылдарының әлеуетін көрсетеді. Нәтижелер гумин қышқылдарының құрылымдық сипаттамалары мен реактивтілігіне геологиялық факторлардың әсері туралы жаңа түсініктер береді, осылайша оларды практикалық қолдану және одан әрі технологиялық дамыту мүмкіндіктерін кеңейтеді. Мұндай айырмашылықтар бұрын бұл кен орындары үшін толық зерттелмеген. Сонымен қатар, зерттеу нәтижелері Кузнецк кен орыны және Күміскұдық кен орындарынан алынған гумин қышқылдарының экологияда, агрохимияда, ауыл шаруашылығында және ағын суларды тазартуда қолдану үшін жоғары әлеуетке ие екенін көрсетті, алайда олардың тиімділігі кен орнына және функционалдық топтардың құрамына байланысты өзгеруі мүмкін. Алынған нәтижелер гумин қышқылдарына негізделген технологияларды дамытуға жаңа мүмкіндіктер ашып, геологиялық факторлардың олардың химиялық құрылымы мен қасиеттеріне әсерін тереңірек түсінуге мүмкіндік береді.

Түйін сөздер: қоңыр көмір, гумин қышқылы, Күміскұдық, Кузнецк кен орны, функционалдық топтар

© С. Тянах^{1*}, Т.О. Хамитова², А.П. Науанова², Д.М. -Кереевна³,
А.С. Дарменбаева⁴, 2025.

¹ Карагандинский университет имени Е.А. Букетова, Караганда, Казахстан;

²Казахский агротехнический исследовательский университет им. Сейфуллина,
Астана, Казахстан;

³Казахского национального университета имени аль-Фараби, Алматы, Казахстан;

⁴Таразский университет имени М.Х. Дулати, Тараз, Казахстан.

E-mail: saika_8989@mail.ru

ИЗУЧЕНИЕ СВОЙСТВ ГУМИНОВЫХ КИСЛОТ, СИНТЕЗИРОВАННЫХ ИЗ БУРЫ УГЛЕЙ КУЗНЕЦКОГО И КУМУСКУДУКСКОГО МЕСТОРОЖДЕНИЙ

Тянах Сайрагул — PhD, Карагандинский университет имени академика Е.А. Букетова, Караганда, Казахстан,

E-mail: saika_8989@mail.ru, <https://orcid.org/0000-0001-5343-4695>;

Хамитова Толкын Ондирисовна — PhD, агрономический факультет, Казахский агротехнический университет им. С. Сейфуллина, Астана, Казахстан,
E-mail: khamitova.t@inbox.ru; <https://orcid.org/0000-0002-4691-3732>;

Науанова Айнаш Пахуашовна — доктор биологических наук, профессор, Казахский агротехнический исследовательский университет им. С. Сейфуллина, Астана, Казахстан,
E-mail: nauanova@mail.ru; <https://orcid.org/0000-0002-5250-1961>;

Дана Мыкты-Кереевна Ибраимова — кандидат химических наук (К.х.н.), старший преподаватель кафедры Аналитической, коллоидной химии и технологии редких элементов химического факультета Казахского национального университета имени аль-Фараби, Алматы, Казахстан,
E-mail: dana_kereevna@kaznu.kz, ORCID: <https://orcid.org/0000-0002-1761-9745>;

Дарменбаева Акмарал Сабетбеккызы — доцент кафедры химии и химической технологии, Таразский университет им. М.Х. Дулати, Тараз, Казахстан,
E-mail: maral88@mail.ru; <https://orcid.org/0000-0003-2974-0398>

Аннотация. Данная статья посвящена синтезу и сравнительному анализу гуминовых кислот, выделенных из бурого угля Кузнецкого и Кумускудукского месторождений. Проводя исследования с использованием современных аналитических методов, таких как инфракрасная спектроскопия, элементный анализ и термогравиметрический анализ, были изучены физико-химические свойства гуминовых кислот, полученных из указанных геологических источников. Исследование выявило значительные различия в химическом составе, функциональных группах и термической стабильности, обусловленные различиями в геологических условиях формирования каждого месторождения. В частности, обнаружены вариации в содержании карбоксильных и гидроксильных групп, которые оказывают влияние на сорбционные и комплексообразующие свойства гуминовых кислот. Проведенное сравнительное исследование подчеркивает перспективность использования гуминовых кислот из обоих месторождений в экологической реабилитации, сельском хозяйстве и очистке сточных вод. Полученные результаты способствуют более глубокому пониманию воздействия геологических факторов на структурные характеристики и реакционную способность гуминовых кислот, открывая новые возможности для их практического применения и дальнейшего технологического совершенствования. Подобные различия ранее подробно не изучались для этих месторождений. Кроме того, исследование показало, что гуминовые кислоты из обоих месторождений имеют высокий потенциал для использования в экологии, агрохимии, сельском хозяйстве и очистке сточных вод, однако их эффективность может различаться в зависимости от месторождения и состава функциональных групп. Полученные результаты открывают новые перспективы для развития технологий на основе гуминовых кислот и позволяют лучше понять влияние геологических факторов на их химическую структуру и свойства.

Ключевые слова: бурый уголь, гуминовые кислоты, Кумускудукское, Кузнецкое месторождение, функциональные группы

Introduction. Kazakhstan ranks among the top ten coal-producing countries in the world, with large deposits of lignite and sub-bituminous coals, especially in regions like Karaganda, Ekibastuz, and the Kumuskuduk fields. These deposits are rich in organic

matter, making them ideal for the extraction of humic acids. The country's vast reserves offer a reliable, cost-effective source of raw material for producing humic substances, positioning Kazakhstan to meet both domestic and international demand. Humic substances are known for their ability to improve soil quality, increase water retention, and remediate polluted soils by binding heavy metals and organic contaminants. Kazakhstan, with its history of industrial activity and mining, faces significant environmental challenges, including soil and water contamination. The extraction and application of humic substances from domestic coal sources offer a promising solution for mitigating environmental degradation, thus stimulating the market for these materials (Polish, 2023; Mendelev, 2024; Tyanakh et al, 2022).

The market for humic substances derived from coals in Kazakhstan is on an upward trajectory, driven by the country's abundant coal resources and the growing demand for sustainable agricultural and environmental solutions. With continued investment and government support, Kazakhstan is well-positioned to expand its production and export capacity, making humic substances a key contributor to its green economy and environmental initiatives (Margaret et al, 2015). Humic acids (HAs) are essential components of natural organic matter, widely distributed in soils, sediments, and water bodies. These complex, heterogeneous macromolecules are formed by the decomposition of plant and animal material over time and play a crucial role in various environmental and biological processes. Humic acids possess unique physicochemical properties, including the ability to chelate metal ions, adsorb organic pollutants, and enhance soil structure. As a result, they are extensively studied for their potential applications in agriculture, environmental remediation, and industry. Despite the significant interest in humic substances, their properties are influenced by their origin and the conditions under which they are formed, necessitating further research on humic acids derived from different natural sources (Chen Y et al, 1976; Kairbekov et al, 2014).

Brown coal, or lignite, represents a promising raw material for the extraction of humic acids due to its high organic content and relatively low degree of carbonization. The Kuznetsk Basin in the central region of Kazakhstan and the Kumuskuduk region are two significant deposits of brown coal with differing geological and environmental characteristics. While these deposits are abundant, the properties of humic acids synthesized from their brown coal remain insufficiently explored. Understanding the relationship between the source material and the properties of the resulting humic acids is critical for optimizing their use in various practical applications (Kairbekov et al, 2023; Dzheldybaeva et al, 2022).

This study focuses on the synthesis of humic acids from brown coal extracted from the Kuznetsk and Kumuskuduk deposits, with the objective of analyzing and comparing their physicochemical properties. Through detailed characterization, including elemental analysis, functional group identification, and molecular weight distribution, this research aims to elucidate the factors that influence the performance of humic acids derived from these distinct coal sources. The findings of this work are expected to contribute to the development of more efficient humic acid-based products and technologies in agriculture, environmental management, and industrial processes,

while also advancing the fundamental understanding of humic substances from different coal deposits (Kairbekov et al, 2014; Arziev et al, 2002; Feng et al, 2024).


Materials and methods. The physico-chemical characteristics of lignite from different deposits can exhibit substantial variability due to the geological conditions under which it formed, as well as the composition of its organic and mineral constituents. Essential parameters for evaluating lignite's suitability for diverse applications—such as energy production, chemical processing, and environmental protection - include moisture content, ash content, volatile matter yield, and the types of functional groups present.

In this study, lignite samples from the Kumuskudyk and Kuznetsk deposits were examined. These samples represent coal resources with specific physicochemical properties, making them valuable for investigating the extraction of humic acids and assessing their potential applications in fields like agriculture and environmental management. Analyzing these samples provides deeper insights into lignite's composition and structure, along with its reactivity with various agents in the process of humic substance extraction.

The Kumuskudyk coal mine, one of Kazakhstan's largest coal-producing sites, is operated by the joint-stock company "Sat Komir Mining Company." This mine, located in the Bukar Zhyrau district of the Karaganda region approximately 35 km east of Karaganda, primarily produces B-3 lignite. Similarly, the Kuznetsk coal mine, an important coal operation in Kazakhstan under the Kuznetsk company, is situated in the same district and approximately the same distance from Karaganda. The Kuznetsk mine is one of the top ten coal producers in Kazakhstan and is also known for its output of B-3 lignite.

In the initial phase of the study, lignite samples from the Kumuskudyk (KUM) and Kuznetsk (KUZ) deposits were selected as raw materials for humic acid extraction. To enhance surface area and improve the efficiency of humic acid extraction, each sample was pre-ground to a particle size of 0.5–1 mm using a "Vibrotechnik ShKD-6" crusher. Subsequently, the coal samples were dried in an oven at 80°C for 8 hours to eliminate moisture.

Extraction of humic acid from coal using the hydro-alkaline method. Humic acids were extracted from lignite by an alkali extraction method, which involves subsequent precipitation of humic acids from alkaline extracts using hydrochloric acid. This method isolates humic acids through their reaction with NaOH, followed by separation via precipitation, as described by the following equation.



Humic acids were extracted from the coal samples by adding NaOH, followed by precipitation using an acidic reagent. During the extraction, 10.0–20.0 g of coal was accurately weighed to 0.0001 g and placed in a 250 cm³ flask. In the next stage, 100

cm³ of a 4% sodium hydroxide solution was added. The mixture was heated to 80°C and stirred on a shaker for 2 hours. The resulting suspension was filtered or centrifuged to remove undissolved carbon residues, which were subsequently rinsed with a small amount of alkaline solution. The remaining solid carbon was dried and weighed for further analysis. The volume of the filtered solution was recorded, and the humic acid content in the final product was determined (Ismailov et al. 2023; Das et al. 2021; Das et al. 2015).

Humic acid was precipitated by adjusting the pH of the solution to 2-3 using a 5% HCl solution. The mixture was then allowed to settle for 60 minutes, after which the humic acid precipitate was collected on a pre-weighed (blue-strip) filter paper. The precipitate was thoroughly rinsed with distilled water to remove any impurities and residual reagents. Following filtration, the filter paper with the precipitate was carefully removed from the Buchner funnel, folded, and pre-dried. It was then placed in a pre-weighed beaker and dried in an oven at 80°C until a constant weight was achieved, ensuring complete moisture removal and enabling accurate measurement of the humic acid mass obtained (Wells et al. 2015; Ulzhalgas, 2022).

The elemental composition of the samples. The elemental composition of the samples was analyzed using a CHNS-O UNICUBE organic elemental analyzer from Elementar Analysensysteme GmbH (Germany). This analysis was carried out in the laboratory of Nazarbayev University. The analyzer operates on the classic Dumas-Pregl method, which involves combustion of the samples in the presence of an oxidizing agent within an inert gas stream. During the combustion process, oxygen was supplied continuously, leading to the formation of analytical gases such as carbon dioxide (CO₂), water (H₂O), molecular nitrogen (N₂), and sulfur dioxide (SO₂). To ensure precise sample weighing, a Mettler Toledo XPR6U Ultra-Microbalance was investigated. Samples were placed in disposable tin boats with a thickness of less than 0.01 mm and sealed with tweezers to prevent sample loss.

Weight range: suitable for use from micro quantities (<1 mg) to macro quantities (approximately 1 g), with a maximum of 15 mg of organic matter. Element concentration ranges are as follows:

- Carbon (C): up to 14 mg absolute or 0–100% (up to 50 mg in CN mode*).
- Hydrogen (H): up to 2 mg absolute or 0–100%.
- Nitrogen (N): up to 10 mg absolute or 0–100%.
- Sulfur (S): up to 3 mg absolute or 0–100%.
- Oxygen (O*): up to 6 mg absolute or 0–100%.

Accuracy: less than 0.1% absolute composition (for a homogeneous substance), depending on sample type, analysis mode, and configuration. Sample combustion was conducted in a quartz reactor within ELEMENTAR's Unicube analyzer, equipped with a high-temperature-resistant ceramic lining, enabling combustion at temperatures up to 1150°C without the use of catalysts. The oxidation column was maintained at 1150°C, while the reduction column was set to 850°C. Each sample was analyzed in triplicate, and the results were averaged for accuracy. Calibration and verification of the equipment were performed using a sulfonamide standard from Elementar Analysensysteme GmbH.

Tin boats with dimensions of 4x4x11 mm and high-purity gases, helium (99.9999%) and oxygen (99.999%), were used in the measurements. The base standard applied was ASTM D5373, which specifies test methods for determining carbon, hydrogen, and nitrogen in coal samples, as well as carbon in coal and coke samples, using combustion analysis techniques.

Results and discussions. Analysis of the elemental composition of samples. The carbon (C), hydrogen (H), nitrogen (N), sulfur (S), and oxygen (O) content in various samples was determined using a CHNS-O UNICUBE elemental analyzer from Elementar Analysensysteme GmbH. The analysis provided average values for the content of oxygen (O), carbon (C), hydrogen (H), nitrogen (N), and sulfur (S) in brown coal from the Kuznetsk open-pit mine (sample KUZ-2024) and in the ash produced after its combustion (sample Zola KUZ-2024). The results are presented in Table 1.

Table 1. Average Content of Oxygen (O), Carbon (C), Hydrogen (H), Nitrogen (N), and Sulfur (S) in Brown Coal from the Kuznetsk Open-Pit Mine and in the Ash Resulting from Its Combustion

№	Name	Oxygen (O), %	Carbon (C), %	Hydrogen (H), %	Nitrogen (N), %	Sulfur (S), %
1	Kuz-2024	24.62	56.02	5.085	0	0.155
2	Zola Kuz-2024	11.56	12.33	0.16	0.43	0

Brown coal from the Kuznetsk open-pit mine is notable for its high carbon content (56.02%), indicating a substantial calorific value and suitability as a fuel source. The elevated oxygen (25.07%) and hydrogen (5.085%) levels suggest a significant amount of volatile compounds, contributing to high reactivity and enhanced combustion efficiency. The absence of nitrogen minimizes the risk of nitrogen oxide (NO_x) formation, which is beneficial for its environmental impact. Additionally, the low sulfur content (0.155%) implies a reduced likelihood of sulfur dioxide (SO_2) emissions, though monitoring SO_2 emissions during combustion remains essential to mitigate air pollution.

The ash sample exhibits a marked reduction in oxygen (to 11.56%), carbon (to 12.33%), and hydrogen (to 0.16%) content compared to the original coal, indicating that a substantial portion of oxygen and hydrogen was released as combustion gases, such as carbon dioxide (CO_2) and water vapor (H_2O). The presence of 12.33% carbon in the ash suggests residual organic material, which could support the use of ash as a component in organomineral fertilizers. The nitrogen content of 0.43% in the ash is likely due to the formation of nitrogen compounds (NO_x) during coal combustion in an air atmosphere.

The reduction in oxygen and carbon content following combustion results from the transformation of these elements into gaseous combustion products. The remaining carbon content in the ash at 12.33% reflects incomplete oxidation, which could allow for the potential reuse of the ash. Hydrogen, initially at 5.085% in the coal, is nearly absent (0.16%) in the ash due to its release as water vapor. Sulfur, initially present at 0.155% in the coal, is entirely absent in the ash, suggesting it volatilized during combustion as gaseous compounds, such as sulfur dioxide (SO_2).

The ash produced from the combustion of brown coal shows potential as a component in organomineral soil ameliorants. The residual carbon and nitrogen content

may contribute to enhancing soil fertility and improving soil structure. The absence of sulfur in the ash is advantageous from an environmental perspective, as excessive sulfur content could result in soil acidification. The low hydrogen and oxygen levels in the ash further indicate its predominantly mineral nature, which can positively influence the soil's physical and chemical properties.

In conclusion, the ash derived from brown coal of the Kuznetsk open-pit mine has potential for use in soil improvement. However, further analysis is required to assess the presence of heavy metals and toxic elements to provide a comprehensive evaluation of its suitability for this purpose.

The elemental analysis of brown coal from the Kumuskuduk open-pit mine revealed that its primary components are carbon (C), oxygen (O), hydrogen (H), nitrogen (N), and sulfur (S). According to the data in Table 1, the average carbon content in these samples was 50.855%, reflecting typical characteristics of low-grade brown coal from this region. This level of carbon suggests a relatively high calorific value, as carbon is the primary energy source in fuel combustion. However, the notable oxygen content (25.07%) indicates a high level of moisture and oxygenated compounds, which diminishes the coal's energy efficiency. The results are presented in Table 2.

Table 2. Average content of oxygen (O), carbon (C), hydrogen (H), nitrogen (N), and sulfur (S) in brown coal from the Kumuskuduk open-pit mine and in the ash produced after its combustion

№	Name	Oxygen (O), %	Carbon (C), %	Hydrogen (H), %	Nitrogen (N), %	Sulfur (S), %
1	Kum-2024	25.07	50.855	5.294	0.455	0.138
2	Zola Kum-2024	11.56	11.14	0.136	0.36	0

The hydrogen content in the sample is 5.294%, which significantly contributes to the calorific value, as hydrogen releases substantial energy upon combustion. Therefore, its presence enhances the fuel's energy potential. The nitrogen content was found to be 0.455%, which is relatively low. This is beneficial for the coal's environmental performance, as higher nitrogen levels can lead to nitrogen oxide (NO_x) emissions during combustion, a source of atmospheric pollution. Additionally, the sulfur content in Kumuskuduk brown coal is 0.138%, which is also low. This low sulfur level makes the coal environmentally preferable, as it minimizes sulfur oxide (SO_x) emissions, reducing the potential for acid rain formation during combustion.

Analysis of the coal ash revealed a notable reduction in carbon content to 11.14%, indicating that a substantial portion of the organic matter was consumed during combustion. Additionally, the minimal levels of hydrogen and sulfur in the ash further confirm their extensive burnout during the combustion process.

The analysis of the elemental composition of brown coal from the Kumuskuduk deposit leads to the following conclusions:

- The coal has a high carbon content (approximately 51%), suggesting a substantial calorific value. However, a relatively high oxygen content (25%) implies significant moisture, potentially reducing the efficiency of its energy use.
- The low sulfur (0.138%) and nitrogen (0.455%) levels contribute to its

environmentally friendly profile, resulting in minimal pollutant emissions during combustion.

- The hydrogen content of 5.294% enhances the coal's calorific value, though moisture and oxygen remain key factors impacting its energy potential.
- Coal ash analysis confirmed an efficient combustion process, evidenced by a high degree of organic burnout and minimal residual elements, such as carbon, hydrogen, and sulfur.

In summary, brown coal from the Kumuskuduk mine shows promise as a fuel source for the energy sector, particularly due to its low sulfur and nitrogen content, which helps to minimize its environmental impact.

The analysis of coal ash composition (sample Zola Kum-2024) suggests its potential for use as a component in organomineral ameliorants. Despite a reduction in carbon content from 50.855% in the original coal to 11.14% in the ash, the remaining carbon in the ash is still significant, indicating the preservation of organic material. This organic component can positively influence soil structure, enhance water retention, and promote the growth of soil microorganisms, which are crucial for improving soil fertility. The nitrogen content (0.36%) in the ash, although low, could contribute to the nitrogen balance of the soil, benefiting plant growth by providing a small but valuable source of this essential nutrient. The absence of sulfur in the ash is beneficial from an environmental perspective, as excessive sulfur content can lead to soil acidification, which harms soil fertility. Thus, the lack of sulfur makes the ash environmentally safe and suitable for use in soil amelioration.

The oxygen content (O) of 11.56% and hydrogen content (H) of 0.136% in the ash indicate the presence of oxidizing components, which may participate in reactions with other substances in the soil. However, these low values suggest that the ash will primarily function as a mineral component, enhancing the physical structure of the soil.

The analytical data for coal ash from the Kumuskuduk coal mine (Zola Kum-2024) suggest its potential as a source of organomineral ameliorants. The ash contains sufficient carbon to enhance soil structure and fertility. Additionally, its low sulfur and nitrogen content renders it suitable for agricultural applications. However, to comprehensively evaluate the ash's agricultural suitability, it is essential to assess the presence of heavy metals and other potentially toxic pollutants. The concentration of these elements could severely restrict the ash's applicability in agronomy and pose risks to both ecosystems and plant health.

Characterization of functional groups. The functional groups in the samples were identified using an FSM 1202 FTIR spectrometer. To prepare the samples, carbon was ground and mixed with potassium bromide (KBr) to form tablets. The spectra were acquired over a wavenumber range of 4000–400 cm^{-1} with a resolution of 4 cm^{-1} . The analysis primarily focused on the absorption bands corresponding to specific functional groups, including carboxyl (-COOH), hydroxyl (-OH), aromatic (C=C), as well as aliphatic hydrocarbon (C-H) groups. The results are illustrated in Figure 1.




Figure 1 - IR spectrum of the sample

As illustrated in Figure 1, the band in the 3800–3600 cm^{-1} region corresponds to water bound to the coal surface or hydroxyl groups formed during chemical reactions. In the 2000–1600 cm^{-1} range, absorption peaks related to C=O bond vibrations are observed, which can be attributed to carbonyl compounds such as ketones, aldehydes, or carboxyl groups. These bands in coal are likely due to oxygen-containing functional groups involved in oxidation processes. The band between 1600–1300 cm^{-1} is associated with the deformation vibrations of C–H and C=C bonds in aromatic rings. Given that coal contains numerous aromatic structures, this range can provide insights into the aromaticity of the carbon skeleton; stronger bands in this region suggest a higher content of aromatic carbons. The 1000–750 cm^{-1} region may indicate the presence of substituted benzene rings (aromatic hydrocarbons) and C–H in-plane vibrations, as well as vibrations of carbon–oxygen bonds in structures such as ethers or alcohols, if these functional groups are present.

Differential scanning calorimetry (TGA)/DSC analysis. Thermogravimetric (TG) and differential scanning calorimetry (DSC) analysis were conducted using a LABSYSTM EvoTG-DTA/DSC instrument (SETARAM, France) to evaluate the thermal stability of the samples and investigate their decomposition behavior upon heating. Coal samples were placed in aluminum oxide crucibles and subjected to a constant heating rate of 10°C/min in an air atmosphere, reaching a maximum temperature of 800°C. During the analysis, changes in sample mass were continuously recorded, allowing for the determination of the temperatures at which key stages of organic matter decomposition occur. This also facilitated the quantification of moisture, volatile matter, fixed carbon, and ash content. Additionally, TGA provided insights into the stability of the organic components of the coal and its potential reactivity. The results are illustrated in Figure 2.




Figure 2 - Results of TG/DSC analysis of Kumuskuduk coal

As shown in Figure 2, the first phase of mass loss occurred between 100–150°C, which is attributed to the evaporation of moisture. The second phase, occurring between 200–400°C, is associated with the release of volatile substances, including low-molecular hydrocarbons and other organic compounds. The third phase, spanning 400–600°C, corresponds to the decomposition of the primary organic matrix of coal, involving the breakdown of complex macromolecules such as humic substances and hydrocarbons. The remaining mass between 600–800°C reflects the mineral content (ash), which was further corroborated by the ash residue observed following the TG analysis.

Characterization of pore sizes and volumes, moisture content, and density. In the course of the experiments, the ash and moisture content of coal samples from the Kumuskuduk (KUM) and Kuznetsk (KUZ) deposits were measured. The results of these analyses are presented in Table 3.

Table 3. Ash and humidity content of the coal samples.

Indicator	KUM	KUZ
Ash content, Aa, %	22	7,4
Humidity, Wa, %	8,24	9,5

The results revealed that the ash content of coal from the Kumuskuduk deposit is 22%, with a moisture content of 8.24%. The volatile substance yield for this sample is 50.86%. In contrast, the coal from the Kuznetsk deposit exhibited a considerably lower ash content of 7.4% and a moisture content of 9.5%. The yield of volatile substances in the Kuznetsk coal samples ranges from 47% to 50%. These findings highlight significant differences in the characteristics of the coals from the two deposits, which could influence the selection of appropriate technologies for their processing and the extraction of humic substances.

Quantitative determination of humic substance yield from coal waste. The experiments resulted in the extraction of humic acids from coal samples of the Kumuskuduk deposit using the alkaline hydrolysis method. From 10 g of Kumuskuduk coal, 0.7 g of humic acids were obtained, corresponding to a yield of 7%. The yield of humic acids for Kuznetsk coal was 6.3%. This made it possible to carry out a more accurate comparison between the two research objects.

In the optimization of humic acid extraction, various process parameters were examined, including alkali solution concentration (1–4%), temperature (20–80°C), and reaction time (30–120 minutes). Experimental results indicated that the most efficient separation of sodium humates occurred with a 4% alkali solution at 80°C for 120 minutes.

Higher alkali concentrations facilitated more complete extraction of humic acids, whereas increasing the temperature above 80°C resulted in a reduction in sodium humate yield (from 4% to 2%) and significant changes in the product composition. This is attributed to the hydrolysis and leaching of carboxyl and polysaccharide fragments, which led to an increased relative content of aromatic structures (up to 44–45%).

Prolonging the reaction time beyond 2 hours did not notably enhance the extraction of humic acids. Therefore, the optimal conditions for humic acid extraction were found to be a 4% alkali solution, 80°C, and a reaction time of 120 minutes.

Spectroscopic analysis of humic substance content and investigation of structural characteristics. Humic acids were extracted from coal samples of the Kumuskuduk and Kuznetsk deposits through alkaline hydrolysis in the course of the experiments. The extracted humic acid samples were subsequently analyzed using infrared (IR) spectroscopy and thermogravimetric (TG) / differential scanning calorimetry (DSC) techniques. The results are illustrated in Figure 3.






Figure 3- Infrared (IR) spectra of the extracted humic acids

As shown in Figure 3, the absorption band in the 3800–3550 cm^{-1} range corresponds to the vibrations of O–H groups, which are characteristic of hydroxyl groups found in both free water and hydrogen bonds within the structure of humic acids. The 2100–

1600 cm^{-1} range displays absorption bands associated with carboxylic acids, aldehydes, ketones, and esters. The 1600–1250 cm^{-1} region is linked to the vibrations of aromatic C=C bonds, typical of humic acids, which contain aromatic rings in their structure. Deformation vibrations of C–O bonds in carboxylic and phenolic groups may also appear in this range, indicating the presence of complex oxygen-containing functional groups in humic acids. Between 1000–600 cm^{-1} , absorption bands are observed for deformation vibrations of C–H bonds in aromatic rings and C–O–C bond vibrations in esters and ether groups. The low-frequency range of 600–500 cm^{-1} corresponds to out-of-plane vibrations in aromatic systems and deformations of C–C bonds in carbon chains. Thermogravimetric analysis of the humic acids was conducted in an inert atmosphere up to a temperature of 1000°C. The results are illustrated in Figure 4.



(a)



(b)

Figure 4 - Results of TG/DSC analysis of humic acid samples (a) - KUZ; (b) – KUM

As observed from the thermogravimetric analysis curves of the humic acids, in the temperature range of 30–175°C, there is only a minor mass change, likely due to the presence of residual water. The decomposition of the sample occurs in two main stages. The first stage, between 200–500°C, is characterized by the breakdown of volatile substances and functional groups such as carboxyl, phenolic, and methoxyl groups. During this stage, small organic molecules associated with aromatic structures are also decomposed, leading to a significant mass loss. This stage involves the release of water, carbon dioxide, and other low-molecular-weight gases. The second stage, occurring between 550–1000°C, involves the degradation of more stable aromatic structures and condensed polycyclic systems, resulting in further mass loss and the formation of carbon residues. This stage is associated with carbonization, where organic matter is converted into carbon, and the formation of a stable residue similar to coke. These stages illustrate the sequential thermal decomposition of humic acids, from the breakdown of lighter functional groups to the formation of stable carbon structures.

Analysis of the chemical composition of the humic product. Subsequently, the elemental composition of carbon (C), hydrogen (H), nitrogen (N), sulfur (S), and oxygen (O) in humic acids (HA) extracted from the brown coals of the Kumuskuduk and Kuznetsk deposits was determined using the CHNS-O UNICUBE elemental analyzer (Elementar Analysensysteme GmbH). This analysis was conducted to assess the elemental composition of the humic acids, providing insights into their chemical properties and potential applications in agriculture.

Table 4 presents the average percentage values of chemical element content in humic acids derived from different deposits.

Table 4. Average content of oxygen (O), carbon (C), hydrogen (H), nitrogen (N), and sulfur (S) in humic acids

№	Name	Carbon (C), %	Hydrogen (H), %	Nitrogen (N), %	Sulfur (S), %	Oxygen (O), %
1	Humic acid (Kuznetsk)	49.605	3.071	0.93	0.391	45.263
2	Humic acid (Kumuskuduk)	46.69	3.079	0.63	0.172	48.26

Humic acid from the Kuznetsk deposit shows a higher carbon content (49.605%) than that from the Kumuskuduk deposit (46.69%), suggesting a greater degree of aromaticity or polymerization in the organic material of Kuznetsk coals. Hydrogen content is nearly identical in both acids, with 3.071% in Kuznetsk and 3.079% in Kumuskuduk, indicating similar hydrogen-containing structural groups in both samples. Nitrogen content differs between the two, with 0.93% in Kuznetsk humic acid and 0.63% in Kumuskuduk, potentially reflecting a higher presence of amino groups or proteins in the Kuznetsk sample. The sulfur content in Kuznetsk humic acid (0.391%) is significantly higher than in Kumuskuduk (0.172%), possibly due to a higher concentration of sulfide or organic sulfur compounds. Kumuskuduk humic acid has a higher oxygen content (48.26%) compared to Kuznetsk (45.263%), which may indicate a greater presence of carboxyl groups, enhancing the acidity and reactivity of the substance.

The elemental analysis results suggest that humic acid from the Kuznetsk deposit

contains higher levels of carbon, nitrogen, and sulfur, which may indicate a more complex structure and potentially distinct properties compared to humic acid from the Kumuskuduk deposit, characterized by a higher oxygen content. These compositional differences likely stem from variations in the coal formation conditions at these deposits and may influence their respective application areas.

Humic acid from the Kuznetsk deposit is the most suitable for producing organomineral ameliorants. Key reasons include:

- **High carbon content (49.605%):** Carbon forms the organic backbone of humic acids, with a high concentration indicating greater aromaticity and polymerization, enhancing the stability of the organic phase in soil. This structure supports long-term soil improvement and boosts fertility.

- **Elevated nitrogen content (0.93%):** Nitrogen is essential for plant nutrition, and its higher concentration in Kuznetsk humic acids enhances their utility in organomineral fertilizers and ameliorants. This nitrogen content promotes plant growth and optimizes soil agrochemical properties.

- **Sulfur content (0.391%):** Although high sulfur can be unfavorable in some cases, it benefits organomineral ameliorants by supporting amino acid and enzyme synthesis crucial for plants, thus enhancing nutrient availability.

In summary, the balanced composition of Kuznetsk humic acid—with its elevated carbon, nitrogen, and sulfur levels—makes it ideal for organomineral ameliorants, improving soil properties and supplying essential nutrients to plants.

Conclusion. The study of lignite from the Kuznetsk and Kumuskuduk deposits yielded valuable insights into their physicochemical characteristics, highlighting their potential for various industrial applications. Key properties, including ash content, moisture, and volatile matter, support the suitability of these coals for use in energy production and industrial processes.

The analysis revealed that lignite from both deposits exhibits low ash content and high volatile matter yield, making it promising for thermal processing techniques such as gasification and pyrolysis. These properties indicate its high potential as a raw material source for the chemical industry, particularly in the production of synthetic gases and liquid hydrocarbons. Elemental analysis, combined with infrared (IR) spectroscopy and thermogravimetric (TG) analysis, provided critical data on the chemical composition and structure of the coals, identifying key functional groups and elemental contents relevant to their industrial potential.

Additionally, humic acids were extracted from coal samples from both deposits using alkaline hydrolysis and further analyzed through elemental analysis, IR spectroscopy, and TG/DSC analysis. The comparative analysis of the physicochemical properties of humic acids from these brown coals revealed substantial differences attributable to the distinct geological formation conditions of the deposits.

Based on the research conducted on humic acids synthesized from brown coals of the Kuznetsk and Kumuskuduk deposits, significant differences in their elemental composition and structural properties were identified. The Kuznetsk-derived humic acid demonstrated higher carbon, nitrogen, and sulfur content, suggesting a more complex molecular structure, with increased aromaticity and polymerization, which

are beneficial for soil stability and fertility enhancement. Conversely, humic acid from the Kumuskuduk deposit exhibited higher oxygen content, potentially contributing to greater reactivity in soil applications.

These compositional differences indicate that humic acid from the Kuznetsk deposit is particularly promising for the production of organomineral ameliorants, as its balanced elemental profile enhances its potential as a soil conditioner and nutrient source. The findings of this study highlight the importance of deposit-specific characteristics in determining the suitability of humic acids for various agricultural and environmental applications. Further research could focus on the application-specific performance of these humic acids in soil ecosystems to validate and optimize their use in agronomy.

Литература

Baikenov M.I., S. Tyanakh, Ma Feng-Yun, Gulmaliev A.M., Makasheva A.M., Khamitova T.O. & Malyshev V.P. (2024) Viscosity model for the middle fraction of Atasu-Alashankou oil sludge, Mendelev Communications. — Vol. 34(3). — P. 446-449. DOI 10.1016/j.mencom.2024.04.043

Chen Y., Banin A., Schnitzer M. (1976) Use of the scanning electron microscope for structural studies on soil and soil components, Scanning electron microscopy. Part 3: Proceeding of the workshop on techniques for particulate matter studies on SEM. — P.425 - 432. (in English)

Das, T., et al. (2015). Characterizations of Humic Acid Isolated from Coals of Two Nagaland, Coalfields of India in Relation to Their Origin. Journal of the Geological Society of India. — Vol. 86(4). — 468-474. DOI:10.1007/s12594-015-0334-0.

Das T., et al. (2021) Coal-Derived Humic Acid for Application in Acid Mine Drainage (AMD) Water Treatment and Electrochemical Devices. International Journal of Coal Science & Technology. — Vol. 8(6). — P. 1479-1490. DOI:10.1007/s40789-021-00441-5

Dzheldybaeva I.M., Kairbekov Z.K., Maloletnev A.S., Abil'mazhinova D.Z., Suimbaeva S.M. (2022) Physicochemical and Antioxidant Properties of Humic Substances from Coals of the Oy-Karagay and Kiyakty Deposits in the Republic of Kazakhstan, Solid Fuel Chemistry. — Vol. 56. — P. 471-477. DOI 10.3103/S0361521921060033

Feng, X., et al. (2024). Preparation of Humic Acid from Weathered Coal by Mechanical Energy, Activation and Its Properties. Minerals. — Vol. 14(7). DOI:10.3390/min14070648.

Ismailov B., et al. (2023). Methods for Obtaining Phosphorus-Containing Fertilizers Based on Industrial Waste. Inorganics. — 2023. — Vol 11(6). — P.224. <https://doi.org/10.3390/inorganics11060224>

Kairbekov Zh.K., Suimbaeva S.M., Dzheldybaeva I.M., Kairbekov A.Zh., Abil'Mazhinova D.Z. (2023) Antioxidant Activity and Bioavailability of Humic Substances of Low-Mineralized Sulphide Mud, Engineered Science. — № 25. — P. 941-948. DOI 10.30919/es941

Kairbekov Zh.K., Toktamysov M.T., Zhalgasuly N., Eshova Zh.T.(2014) Kompleksnaya pererabotka burykh uglei Tsentral'nogo Kazakhstana (Integrated Processing of Brown Coal from Central Kazakhstan), Almaty: Izd. KazNU..

Margaret Suárez Muñoz, Clara Melián Rodríguez, Alina Gelen (2015) Rudnikas and other autors. Physicochemical characterization, elemental speciation and hydrogeochemical modeling of river and peloid sediments used for therapeutic uses, Applied Clay Science. — Vol. 104. — P. 36-47. DOI 10.1016/j.clay.2014.11.029

Tyanakh S., Baikenov M.I., Gulmaliev A.M., Ma, Feng-Yun, Musina G., Khamitova T.O., & Bolatbay A.N. (2022) Kinetics of Thermolysis of a Low-Temperature Tar in the Presence of a Catalyzer Agent with Deposited Metals, Bulletin of the University of Karaganda Chemistry. — № 4(108). — P. 89-98. DOI 10.31489/2022Ch4/4-22-19

Tyanakh S., Baikenov M.I., Ma Feng Yun, Khamitova T.O., Balpanova N.Zh., Tulebayeva B., Kyzkenova A., Karimova A.B., Rakhimzhanova N.Zh., Kochegina E.V. (2023) Kinetic of oil sludge thermolysis process in presence of nickel, cobalt and iron-supported microsilicate, Polish Journal of Chemical Technology. — Vol. 23(3). — P.101-109. DOI 10.2478/pjct-2023-0030

Ulzhalgas Nazarbek, Perizat Abdurazova, Yerkebulan Raiymbekov (2022) Extraction and Characterization of Humic Acid Based on Coal Mining Waste, *Chem. Eng. Technol.* — V. 45. — No. 6. — P.1133–1140 DOI: 10.1002/ceat.202200038 8(6), 1479-1490. DOI:10.1007/s40789-021-00441-5.

Wells L., Smith, T. J. (2015) Statistical Methods for Chemists: An Introduction to Data Analysis. *Journal of Chemical Education.* — Vol. 92. — no. 6. — P. 1051-1058.

Арзиев Ж.А. (2002) Угли Кыргызстана как основа для получения гуминовых удобрений и стимуляторов роста растений, *Современные проблемы науки и техники: Сб. научн. тр. региональной научн. - теорет. конф., Жалалабатский гос.техн.ин-т.- Жалалабат.* — Ч. 1. — С. 34-38.

Каирбеков Ж.К. (2014) *Переработка твердых горючих ископаемых, Книга, - Алматы: типография «BTS print».* — 260 с.

References

Baikenov M.I., S. Tyanakh, Ma Feng-Yun, Gulmaliev A.M., Makasheva A.M., Khamitova T.O. & Malyshev V.P. (2024) Viscosity model for the middle fraction of Atasu-Alashankou oil sludge, *Mendelev Communications.* — Vol. 34(3). — P.446-449. DOI 10.1016/j.mencom.2024.04.043 (in English)

Chen Y., Banin A., Schnitzer M. (1976) Use of the scanning electron microscope for structural studies on soil and soil components, *Scanning electron microscopy. Part 3: Proceeding of the workshop on techniques for particulate matter studies on SEM.* — P.425 - 432.

Das, T., et al. (2015). Characterizations of Humic Acid Isolated from Coals of Two Nagaland, Coalfields of India in Relation to Their Origin. *Journal of the Geological Society of India.* — Vol. 86(4). — 468-474. DOI:10.1007/s12594-015-0334-0. (in English)

Das T., et al. (2021) Coal-Derived Humic Acid for Application in Acid Mine Drainage (AMD) Water Treatment and Electrochemical Devices. *International Journal of Coal Science & Technology.* — Vol. 8(6). — P. 1479-1490. DOI:10.1007/s40789-021-00441-5 (in English)

Dzheldybaeva I.M., Kairbekov Z.K., Maloletnev A.S., Abil'mazhinova D.Z., Suimbaeva S.M. (2022) Physicochemical and Antioxidant Properties of Humic Substances from Coals of the Oy-Karagay and Kiyakty Deposits in the Republic of Kazakhstan, *Solid Fuel Chemistry.* — Vol. 56. — P. 471-477. DOI 10.3103/S0361521921060033 (in English)

Feng, X., et al. (2024). Preparation of Humic Acid from Weathered Coal by Mechanical Energy, Activation and Its Properties. *Minerals.* — Vol. 14(7). DOI:10.3390/min14070648 (in English)

Ismailov B., et al. (2023). Methods for Obtaining Phosphorus-Containing Fertilizers Based on Industrial, Waste. *Inorganics.* — 2023. — Vol 11(6). — P.224. <https://doi.org/10.3390/inorganics11060224> (in English)

Kairbekov Zh.K., Suimbaeva S.M., Dzheldybaeva I.M., Kairbekov A.Zh., Abil'Mazhinova D.Z. (2023) Antioxidant Activity and Bioavailability of Humic Substances of Low-Mineralized Sulphide Mud, *Engineered Science.* — № 25. — P. 941-948. DOI 10.30919/es941 (in English)

Kairbekov Zh.K., Toktamysov M.T., Zhalgasuly N., Eshova Zh.T.(2014) *Kompleksnaya pererabotka burykh uglei Tsentral'nogo Kazakhstana (Integrated Processing of Brown Coal from Central Kazakhstan),* Almaty: Izd. KazNU. (in English)

Margaret Suárez Muñoz, Clara Melián Rodríguez, Alina Gelen (2015) Rudnikas and other autors. Physicochemical characterization, elemental speciation and hydrogeochemical modeling of river and peloid sediments used for therapeutic uses, *Applied Clay Science.* — Vol. 104. — P. 36-47. DOI 10.1016/j.clay.2014.11.029 (in English)

Tyanakh S., Baikenov M.I., Gulmaliev A.M., Ma, Feng-Yun, Musina G., Khamitova T.O., & Bolatbay A.N. (2022) Kinetics of Thermolysis of a Low-Temperature Tar in the Presence of a Catalyzer Agent with Deposited Metals, *Bulletin of the University of Karaganda Chemistry.* — № 4(108). — P. 89-98. DOI 10.31489/2022Ch4/4-22-19 (in English)

Tyanakh S., Baikenov M.I., Ma Feng Yun, Khamitova T.O., Balpanova N.Zh., Tulebayeva B., Kyzkenova A., Karimova A.B., Rakhimzhanova N.Zh., Kochegina E.V. (2023) Kinetic of oil sludge thermolysis process in presence of nickel, cobalt and iron-supported microsilicate, *Polish Journal of Chemical Technology.* — Vol. 23(3). — .101-109. DOI 10.2478/pjct-2023-0030 (in English)

Ulzhalgas Nazarbek, Perizat Abdurazova, Yerkebulan Raiymbekov (2022) Extraction and

Characterization of Humic Acid Based on Coal Mining Waste, Chem. Eng. Technol. — V. 45. — No. 6. — P.1133–1140 DOI: 10.1002/ceat.202200038 8(6), 1479-1490. DOI:10.1007/s40789-021-00441-5 (in English)

Wells L., Smith, T. J. (2015) Statistical Methods for Chemists: An Introduction to Data Analysis. Journal of Chemical Education. — Vol. 92. —no. 6. — P. 1051-1058 (in English)

Arziev Zh.A. Ugli Kyrgyzstana kak osnova dlja poluchenija guminovyh udobrenij i stimulatorov rosta rastenij, Sovremennye problemy nauki i tehniki: Sb. nauchn. tr. regional'noj nauchn. - teoret. Konf, Zhalalabatskij gos.tehn.in-t [Coals of Kyrgyzstan as a Basis for the Production of Humic Fertilizers and Plant Growth Stimulators, Modern Problems of Science and Technology: Proceedings of the Regional Scientific and Theoretical Conference, Jalal-Abad State Technical Institute.]. Zhalalabat, 2002. — № 1. — P.34 - 38. (in Russian)

Kairbekov Zh.K. (2014). Pererabotka tverdyh gorjuchih iskopaemyh, Kniga [Processing of Solid Fossil Fuels]. — Almaty: tipografija «BTS print», 2014. — P. 260] (in Russian)

Publication Ethics and Publication Malpractice in the journals of the Central Asian Academic Research Center LLP

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the journals of the Central Asian Academic Research Center LLP implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The Central Asian Academic Research Center LLP follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the Central Asian Academic Research Center LLP.

The Editorial Board of the Central Asian Academic Research Center LLP will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Ответственный редактор *А. Ботанқызы*

Редакторы: *Д.С. Аленов, Т. Апендиев*

Верстка на компьютере *Г.Д. Жадырановой*

Подписано в печать 29.09.2025.

Формат 60x88¹/₈. 18,0 п.л.

Заказ 3.

«Central Asian Academic Research Center» LLP

Алматы, Қонаев к-сі, 142