

**ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

ӘЛЬ-ФАРАБИ АТЫНДАҒЫ
ҚАЗАҚ ҰЛТТЫҚ УНИВЕРСИТЕТИНІҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

КАЗАХСКИЙ НАЦИОНАЛЬНЫЙ
УНИВЕРСИТЕТ ИМЕНИ АЛЬ-ФАРАБИ

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

AL-FARABI KAZAKH
NATIONAL UNIVERSITY

ФИЗИКА-МАТЕМАТИКА СЕРИЯСЫ

СЕРИЯ ФИЗИКО-МАТЕМАТИЧЕСКАЯ

PHYSICO-MATHEMATICAL SERIES

5 (321)

ҚЫРКҮЙЕК – ҚАЗАН 2018 ж.
СЕНТЯБРЬ – ОКТЯБРЬ 2018 г.
SEPTEMBER – OCTOBER 2018

1963 ЖЫЛДЫН ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

Бас редакторы
ф.-м.ғ.д., проф., КР ҮФА академигі **F.M. Мұтанов**

Редакция алқасы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев Ү.Ү. проф. корр.-мүшесі (Қазақстан)
Жусіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Сұраған Ә. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«КР ҮФА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» РКБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы қуәлік

Мерзімділігі: жылдана 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / physics-mathematics.kz

© Қазақстан Республикасының Үлттық ғылым академиясы, 2018

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Р е д а к ц и о н на я кол л е г и я:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / physics-mathematics.kz

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskyi I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
www.nauka-nanrk.kz / physics-mathematics.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2018.2518-1726.5>

Volume 5, Number 321 (2018), 33 – 41

UDK 517.951
 MRNTI 27.31.15

A.T.Assanova¹, B.Zh.Alikhanova², K.Zh.Nazarova²

¹ Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;
² Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan
 E-mail: assanova@math.kz; bota_mktu@mail.ru; ginazarova@mail.ru

**WELL-POSEDNESS OF A NONLOCAL PROBLEM
 WITH INTEGRAL CONDITIONS FOR THIRD ORDER
 SYSTEM OF THE PARTIAL DIFFERENTIAL EQUATIONS**

Abstract. The nonlocal problem with integral conditions for the system of partial differential equations third-order is considered. The existence and uniqueness of classical solution to nonlocal problem with integral conditions for third-order system of partial differential equations are studied and the method for constructing their approximate solutions is proposed. Conditions of an unique solvability to nonlocal problem with integral conditions for third order system of partial differential equations are established. By introduction of new unknown functions, we have reduced the considered problem to an equivalent problem consisting of a nonlocal problem with integral conditions and parameters for a system of hyperbolic equations of second order and a integral relations. We have offered the algorithm for finding approximate solution to investigated problem and have proved its convergence. Sufficient conditions for the existence of unique solution to the equivalent problem with parameters are obtained. Well-posedness of the nonlocal problem with integral conditions for third order system of partial differential equations are established in the terms of well-posedness to nonlocal problem with integral conditions for system of hyperbolic equations second order.

Key Words: third order partial differential equations, nonlocal problem, integral condition, system of hyperbolic equations second order, solvability, algorithm.

1. Introduction. In recent years, there has been a great interest to a nonlocal problems for third order partial differential equations and systems. Such problems are appeared in the mathematical modeling of various natural science processes [1-9]. A lot of many works devoted to the investigate of various problems for third order partial differential equations with two independent variables, bibliography and analysis can be see in [1, 2, 5, 7-9]. The third order system of partial differential equations began to be studied relatively recently [5, 7-9].

In the present paper we consider the nonlocal problem with integral conditions for third order system of partial differential equations at a rectangular domain. We investigate the questions of existence and uniqueness of the classical solution to nonlocal problem with integral conditions for third order system of partial differential equations and its applications. Some types of initial-boundary value problems for third order system of partial differential equations are considered in [7-9].

Methods. For solve to considered problem we use a method of introduction additional functional parameters [10-32]. The original problem is reduced to an equivalent problem consisting from nonlocal problem with integral conditions for system of hyperbolic equations second order, containing functional parameters and integral relations. Sufficient conditions of the unique solvability to investigated problem are established in the terms of unique solvability to nonlocal problem with integral conditions for system of hyperbolic equations. Algorithms for finding of approximate solution to the equivalent problem are

constructed. Well-posedness to nonlocal problem with integral conditions for third order system of partial differential equations are established in the terms of well-posedness to nonlocal problem with integral conditions for system of hyperbolic equations second order.

2. *Statement of problem.* At the domain $\Omega = [0, T] \times [0, \omega]$ we consider the following initial-boundary value problem for the special system of partial differential equations

$$\frac{\partial^3 u}{\partial t \partial x^2} = A(t, x) \frac{\partial^2 u}{\partial x^2} + B(t, x) \frac{\partial^2 u}{\partial t \partial x} + C(t, x) \frac{\partial u}{\partial x} + D(t, x) \frac{\partial u}{\partial t} + E(t, x)u + f(t, x), \quad (t, x) \in \Omega, \quad (1)$$

$$\int_0^T L(\tau, x) \frac{\partial u(\tau, x)}{\partial x} d\tau = \varphi(x), \quad x \in [0, \omega], \quad (2)$$

$$u(t, 0) = \psi_0(t), \quad t \in [0, T], \quad (3)$$

$$\int_0^\omega P(t, \xi) \frac{\partial u(t, \xi)}{\partial \xi} d\xi = \psi_1(t), \quad t \in [0, T], \quad (4)$$

where $u(t, x) = \text{col}(u_1(t, x), u_2(t, x), \dots, u_n(t, x))$ is unknown function, the $n \times n$ -matrices $A(t, x)$, $B(t, x)$, $C(t, x)$, $D(t, x)$, $E(t, x)$ and n -vector function $f(t, x)$ are continuous on Ω , the $n \times n$ -matrix $L(t, x)$ is continuous and continuously differentiable by x on Ω , the n -vector-function $\varphi(x)$ is continuously differentiable on $[0, \omega]$, the $n \times n$ -matrix $P(t, x)$ is continuous and continuously differentiable by t on Ω , the n -vector-functions $\psi_0(t)$ and $\psi_1(t)$ are continuously differentiable on $[0, T]$.

A function $u(t, x) \in C(\Omega, R^n)$ having partial derivatives $\frac{\partial u(t, x)}{\partial x} \in C(\Omega, R^n)$, $\frac{\partial u(t, x)}{\partial t} \in C(\Omega, R^n)$, $\frac{\partial^2 u(t, x)}{\partial x^2} \in C(\Omega, R^n)$, $\frac{\partial^2 u(t, x)}{\partial t \partial x} \in C(\Omega, R^n)$, $\frac{\partial^3 u(t, x)}{\partial t \partial x^2} \in C(\Omega, R^n)$ is called a classical solution to problem (1)-(4) if it satisfies system (1) for all $(t, x) \in \Omega$, and integral and initial conditions (2), (3) and (4).

We investigate the questions of existence and uniqueness of the classical solutions to nonlocal problem with integral conditions for system of partial differential equations of third order (1)-(4) and the approaches of constructing its approximate solutions. For this goals, we applied the method of introduction additional functional parameters proposed in [10-32] for the solve of nonlocal boundary value problems for systems of hyperbolic equations with mixed derivatives. Considered problem is provided to nonlocal problem with integral conditions for system of hyperbolic equations of second order, including additional functions and integral relations. The algorithm of finding the approximate solution of the investigated problem is proposed and its convergence proved. Sufficient conditions of the existence unique classical solution to problem (1)-(4) are obtained in the terms of unique solvability to nonlocal problem with integral conditions for system of hyperbolic equations second order.

3. Scheme of the method and reduction to equivalent problem.

We introduce a new unknown function $v(t, x) = \frac{\partial u(t, x)}{\partial x}$ and re-write nonlocal problem with integral conditions (1)-(4) in the following form

$$\frac{\partial^2 v}{\partial t \partial x} = A(t, x) \frac{\partial v}{\partial x} + B(t, x) \frac{\partial v}{\partial t} + C(t, x)v + D(t, x) \frac{\partial u}{\partial t} + E(t, x)u + f(t, x), \quad (t, x) \in \Omega, \quad (5)$$

$$\int_0^T L(\tau, x)v(\tau, x)d\tau = \varphi(x), \quad x \in [0, \omega], \quad (6)$$

$$\int_0^\omega P(t, \xi) v(t, \xi) d\xi = \psi_1(t), \quad t \in [0, T], \quad (7)$$

$$u(t, x) = \psi_0(t) + \int_0^x v(t, \xi) d\xi, \quad \frac{\partial u(t, x)}{\partial t} = \dot{\psi}_0(t) + \int_0^x \frac{\partial v(t, \xi)}{\partial t} d\xi, \quad (t, x) \in \Omega. \quad (8)$$

Here the condition (3) is taken account in (8).

A pair functions $(v(t, x), u(t, x))$, where the function $v(t, x) \in C(\Omega, R^n)$ has partial derivatives $\frac{\partial v(t, x)}{\partial x} \in C(\Omega, R^n)$, $\frac{\partial v(t, x)}{\partial t} \in C(\Omega, R^n)$, $\frac{\partial^2 v(t, x)}{\partial t \partial x} \in C(\Omega, R^n)$, the function $u(t, x) \in C(\Omega, R^n)$ has partial derivatives $\frac{\partial u(t, x)}{\partial x} \in C(\Omega, R^n)$, $\frac{\partial u(t, x)}{\partial t} \in C(\Omega, R^n)$, $\frac{\partial^2 u(t, x)}{\partial x^2} \in C(\Omega, R^n)$, $\frac{\partial^2 u(t, x)}{\partial t \partial x} \in C(\Omega, R^n)$, $\frac{\partial^3 u(t, x)}{\partial t \partial x^2} \in C(\Omega, R^n)$, is called a solution to problem (5)–(8) if it satisfies of the system of hyperbolic equations (5) for all $(t, x) \in \Omega$, the boundary conditions (6), (7), and the integral relation (8).

At fixed $u(t, x)$ the problem (5)–(7) is the nonlocal problem with integral conditions for the system of hyperbolic equations with respect to $v(t, x)$ on Ω . The integral relations (8) allow us to determine the unknown functions $u(t, x)$ and its partial derivative $\frac{\partial u(t, x)}{\partial t}$ for all $(t, x) \in \Omega$.

4. Algorithm for finding of the approximate solution to problem (5)–(8).

The unknown function $v(t, x)$ will be determined from nonlocal problem with integral conditions for system of hyperbolic equations (5)–(7). The unknown function $u(t, x)$ and its partial derivative $\frac{\partial u(t, x)}{\partial t}$ will be found from integral relations (8). If we known the function $u(t, x)$ and its partial derivative $\frac{\partial u(t, x)}{\partial t}$, then from nonlocal problem with integral conditions (5)–(7) we find the function $v(t, x)$.

Conversely, if we known the function $v(t, x)$ and its partial derivative $\frac{\partial v(t, x)}{\partial t}$, then from integral relations (8) we find the function $u(t, x)$ and its partial derivative $\frac{\partial u(t, x)}{\partial t}$. Since the functions $u(t, x)$ and $v(t, x)$ are unknowns together for finding of the solution to problem (5)–(8) we use an iterative method.

The solution to problem (5)–(8) is the pair functions $(v^*(t, x), u^*(t, x))$ we defined as a limit of sequence of pairs $(v^{(k)}(t, x), u^{(k)}(t, x))$, $k = 0, 1, 2, \dots$, according to the following algorithm:

- Step 0.* 1) Suppose in the right-hand part of system (5) $u(t, x) = \psi_0(t)$, $\frac{\partial u(t, x)}{\partial t} = \dot{\psi}_0(t)$, from nonlocal problem with integral conditions (5)–(7) we find the initial approximation $v^{(0)}(t, x)$ and its partial derivatives for all $(t, x) \in \Omega$;
- 2) From integral relations (8) under $v(t, x) = v^{(0)}(t, x)$, $\frac{\partial v(t, x)}{\partial t} = \frac{\partial v^{(0)}(t, x)}{\partial t}$, we find the functions $u^{(0)}(t, x)$ and $\frac{\partial u^{(0)}(t, x)}{\partial t}$ for all $(t, x) \in \Omega$.

Step 1. 1) Suppose in the right-hand part of system (5) $u(t, x) = u^{(0)}(t, x)$, $\frac{\partial u(t, x)}{\partial t} = \frac{\partial u^{(0)}(t, x)}{\partial t}$,

from nonlocal problem with integral conditions (5)-(7) we find the first approximation $v^{(1)}(t, x)$ and its partial derivatives for all $(t, x) \in \Omega$.

2) From integral relations (8) under $v(t, x) = v^{(1)}(t, x)$, $\frac{\partial v(t, x)}{\partial t} = \frac{\partial v^{(1)}(t, x)}{\partial t}$, we find the functions $u^{(1)}(t, x)$ and $\frac{\partial u^{(1)}(t, x)}{\partial t}$ for all $(t, x) \in \Omega$.

And so on.

Step k . 1) Suppose in the right-hand part of system (5) $u(t, x) = u^{(k-1)}(t, x)$, $\frac{\partial u(t, x)}{\partial t} = \frac{\partial u^{(k-1)}(t, x)}{\partial t}$, from nonlocal problem with integral conditions (5)-(7) we find the k -th approximation $v^{(k)}(t, x)$ and its partial derivatives for all $(t, x) \in \Omega$:

$$\frac{\partial^2 v^{(k)}}{\partial t \partial x} = A(t, x) \frac{\partial v^{(k)}}{\partial x} + B(t, x) \frac{\partial v^{(k)}}{\partial t} + C(t, x) v^{(k)} + D(t, x) \frac{\partial u^{(k-1)}}{\partial t} + E(t, x) u^{(k-1)} + f(t, x), \quad (t, x) \in \Omega, \quad (9)$$

$$\int_0^T L(\tau, x) v^{(k)}(\tau, x) d\tau = \varphi(x), \quad x \in [0, \omega], \quad (10)$$

$$\int_0^\omega P(t, \xi) v^{(k)}(t, \xi) d\xi = \psi_1(t), \quad t \in [0, T]. \quad (11)$$

2) From integral relations (8) under $v(t, x) = v^{(k)}(t, x)$, $\frac{\partial v(t, x)}{\partial t} = \frac{\partial v^{(k)}(t, x)}{\partial t}$, we find the function $u^{(k)}(t, x)$ and $\frac{\partial u^{(k)}(t, x)}{\partial t}$ for all $(t, x) \in \Omega$:

$$u^{(k)}(t, x) = \psi_0(t) + \int_0^x v^{(k)}(t, \xi) d\xi, \quad \frac{\partial u^{(k)}(t, x)}{\partial t} = \dot{\psi}_0(t) + \int_0^x \frac{\partial v^{(k)}(t, \xi)}{\partial t} d\xi, \quad (t, x) \in \Omega. \quad (12)$$

$k = 1, 2, 3, \dots$

5. The main result.

Consider nonlocal problem with integral conditions (5)-(7) at fixed $u(t, x)$. Then nonlocal problem with integral conditions for system of hyperbolic equations can be have the following form

$$\frac{\partial^2 v}{\partial t \partial x} = A(t, x) \frac{\partial v}{\partial x} + B(t, x) \frac{\partial v}{\partial t} + C(t, x) v + F(t, x), \quad (t, x) \in \Omega, \quad (13)$$

$$\int_0^T L(\tau, x) v(\tau, x) d\tau = \varphi(x), \quad x \in [0, \omega], \quad (14)$$

$$\int_0^\omega P(t, \xi) v(t, \xi) d\xi = \psi_1(t), \quad t \in [0, T], \quad (15)$$

where n -vector function $F(t, x)$ is continuous on Ω .

The following theorem gives conditions of feasibility and convergence of the constructed algorithm and the conditions of the existence unique solution to problem (5)-(8).

Theorem 1. Suppose that

i) the $n \times n$ -matrices $A(t, x)$, $B(t, x)$, $C(t, x)$, $D(t, x)$, $E(t, x)$, and n -vector function $f(t, x)$

are continuous on Ω ;

ii) the $n \times n$ -matrix $L(t, x)$ is continuous and continuously differentiable by x on Ω ; and the n -vector-function $\varphi(x)$ continuously differentiable on $[0, \omega]$;

iii) the $n \times n$ -matrix $P(t, x)$ is continuous and continuously differentiable by t on Ω ; the n -vector-functions $\psi_0(t)$ and $\psi_1(t)$ are continuously differentiable on $[0, T]$;

iv) Nonlocal problem with integral conditions for system of hyperbolic equations (13)–(15) has a unique classical solution.

Then equivalent nonlocal problem for system of hyperbolic equations with integral conditions and parameters (5)–(8) has a unique solution.

Proof. Let the conditions i) – iv) of Theorem 1 are fulfilled. By the algorithm, proposing above, on 0th step we have

$$\frac{\partial^2 v^{(0)}}{\partial t \partial x} = A(t, x) \frac{\partial v^{(0)}}{\partial x} + B(t, x) \frac{\partial v^{(0)}}{\partial t} + C(t, x)v^{(0)} + D(t, x)\dot{\psi}_0(t) + E(t, x)\psi_0(t) + f(t, x), \quad (t, x) \in \Omega, \quad (16)$$

$$\int_0^T L(\tau, x)v^{(0)}(\tau, x)d\tau = \varphi(x), \quad x \in [0, \omega], \quad (17)$$

$$\int_0^\omega P(t, \xi)v^{(0)}(t, \xi)d\xi = \psi_1(t), \quad t \in [0, T]. \quad (18)$$

Since condition iv) is valid, problem (16)–(18) has a unique classical solution $v^{(0)}(t, x)$ and the following inequality holds

$$\begin{aligned} & \max \left(\max_{(t,x) \in \Omega} \|v^{(0)}(t, x)\|, \max_{(t,x) \in \Omega} \left\| \frac{\partial v^{(0)}(t, x)}{\partial x} \right\|, \max_{(t,x) \in \Omega} \left\| \frac{\partial v^{(0)}(t, x)}{\partial t} \right\| \right) \leq \\ & \leq K(x) \max \left(\max_{(t,x) \in \Omega} \|f(t, x)\|, \max_{t \in [0, T]} \|\psi_1(t)\|, \max_{x \in [0, \omega]} \|\varphi(x)\|, \max_{(t,x) \in \Omega} \|F^{(0)}(t, x)\| \right), \end{aligned}$$

where the function $K(x)$ is continuous on $[0, \omega]$, positive and independent on functions $f(t, x)$, $\psi_1(t)$, $\varphi(x)$, $F^{(0)}(t, x) = D(t, x)\dot{\psi}_0(t) + E(t, x)\psi_0(t)$.

Initial approximations $u^{(0)}(t, x)$ and $\frac{\partial u^{(0)}(t, x)}{\partial t}$ are determined by the following form

$$u^{(0)}(t, x) = \psi_0(t) + \int_0^x v^{(0)}(t, \xi)d\xi, \quad \frac{\partial u^{(0)}(t, x)}{\partial t} = \dot{\psi}_0(t) + \int_0^x \frac{\partial v^{(0)}(t, \xi)}{\partial t}d\xi, \quad (t, x) \in \Omega. \quad (19)$$

Then functions $u^{(0)}(t, x)$ and $\frac{\partial u^{(0)}(t, x)}{\partial t}$ satisfy the estimate

$$\begin{aligned} & \max \left(\max_{t \in [0, T]} \|u^{(0)}(t, x)\|, \max_{t \in [0, T]} \left\| \frac{\partial u^{(0)}(t, x)}{\partial t} \right\| \right) \leq \max \left(\max_{t \in [0, T]} \|\psi_0(t)\|, \max_{t \in [0, T]} \|\dot{\psi}_0(t)\| \right) + \\ & + \int_0^x \max \left(\max_{t \in [0, T]} \|v^{(0)}(t, \xi)\|, \max_{t \in [0, T]} \left\| \frac{\partial v^{(0)}(t, \xi)}{\partial t} \right\| \right) d\xi. \end{aligned}$$

For k th approximations determined by relations (9)–(12), we have

$$\max \left(\max_{(t,x) \in \Omega} \|v^{(k)}(t, x)\|, \max_{(t,x) \in \Omega} \left\| \frac{\partial v^{(k)}(t, x)}{\partial x} \right\|, \max_{(t,x) \in \Omega} \left\| \frac{\partial v^{(k)}(t, x)}{\partial t} \right\| \right) \leq$$

$$\leq K(x) \max \left(\max_{(t,x) \in \Omega} \|f(t,x)\|, \max_{t \in [0,T]} \|\psi_1(t)\|, \max_{x \in [0,\omega]} \|\varphi(x)\|, \max_{(t,x) \in \Omega} \|F^{(k-1)}(t,x)\| \right), \quad (20)$$

$$\begin{aligned} & \max \left(\max_{t \in [0,T]} \|u^{(k)}(t,x)\|, \max_{t \in [0,T]} \left\| \frac{\partial u^{(k)}(t,x)}{\partial t} \right\| \right) \leq \max \left(\max_{t \in [0,T]} \|\psi_0(t)\|, \max_{t \in [0,T]} \|\dot{\psi}_0(t)\| \right) + \\ & + \int_0^x \max \left(\max_{t \in [0,T]} \|v^{(k)}(t,\xi)\|, \max_{t \in [0,T]} \left\| \frac{\partial v^{(k)}(t,\xi)}{\partial t} \right\| \right) d\xi, \end{aligned} \quad (21)$$

where $F^{(k-1)}(t,x) = D(t,x) \frac{\partial u^{(k-1)}(t,x)}{\partial t} + E(t,x)u^{(k-1)}(t,x)$.

$$\begin{aligned} \text{Let } \Delta v^{(k)}(t,x) &= v^{(k+1)}(t,x) - v^{(k)}(t,x), \quad \Delta_x v^{(k)}(t,x) = \frac{\partial v^{(k+1)}(t,x)}{\partial x} - \frac{\partial v^{(k)}(t,x)}{\partial x}, \\ \Delta_t v^{(k)}(t,x) &= \frac{\partial v^{(k+1)}(t,x)}{\partial t} - \frac{\partial v^{(k)}(t,x)}{\partial t}, \quad \Delta u^{(k)}(t,x) = u^{(k+1)}(t,x) - u^{(k)}(t,x), \\ \Delta_t u^{(k)}(t,x) &= \frac{\partial u^{(k+1)}(t,x)}{\partial t} - \frac{\partial u^{(k)}(t,x)}{\partial t}, \quad k = 1, 2, 3, \dots. \end{aligned}$$

Then, for differences $\Delta v^{(k)}(t,x)$, $\Delta_x v^{(k)}(t,x)$, $\Delta_t v^{(k)}(t,x)$, $\Delta u^{(k)}(t,x)$, $\Delta_t u^{(k)}(t,x)$ are valid the following inequalities

$$\begin{aligned} & \max \left(\max_{(t,x) \in \Omega} \|\Delta v^{(k)}(t,x)\|, \max_{(t,x) \in \Omega} \|\Delta_x v^{(k)}(t,x)\|, \max_{(t,x) \in \Omega} \|\Delta_t v^{(k)}(t,x)\| \right) \leq \\ & \leq K(x) \max \left(\max_{(t,x) \in \Omega} \|\Delta u^{(k-1)}(t,x)\|, \max_{(t,x) \in \Omega} \|\Delta_t u^{(k-1)}(t,x)\| \right), \end{aligned} \quad (22)$$

$$\max \left(\max_{t \in [0,T]} \|\Delta u^{(k)}(t,x)\|, \max_{t \in [0,T]} \|\Delta_t u^{(k)}(t,x)\| \right) \leq \int_0^x \max \left(\max_{t \in [0,T]} \|\Delta v^{(k)}(t,\xi)\|, \max_{t \in [0,T]} \|\Delta_t v^{(k)}(t,\xi)\| \right) d\xi. \quad (23)$$

From estimates (20)–(23) is follows

$$\begin{aligned} \max \left(\max_{t \in [0,T]} \|\Delta u^{(k)}(t,x)\|, \max_{t \in [0,T]} \|\Delta_t u^{(k)}(t,x)\| \right) &\leq \frac{[\hat{K}x]^k}{k!} \cdot \max \left(\max_{(t,x) \in \Omega} \|\Delta v^{(0)}(t,x)\|, \max_{(t,x) \in \Omega} \|\Delta_t v^{(0)}(t,x)\| \right), \\ \text{where } \hat{K} &= \max_{x \in [0,\omega]} K(x). \end{aligned}$$

Hence, we obtain the uniform convergence of the sequences $\{u^{(k)}(t,x)\}$ and $\left\{ \frac{\partial u^{(k)}(t,x)}{\partial t} \right\}$ to functions $u^*(t,x)$ and $\frac{\partial u^*(t,x)}{\partial t}$ on Ω , respectively, as $k \rightarrow \infty$.

Then the sequences $\{v^{(k)}(t,x)\}$, $\left\{ \frac{\partial v^{(k)}(t,x)}{\partial x} \right\}$, and $\left\{ \frac{\partial v^{(k)}(t,x)}{\partial t} \right\}$ also will be convergent to functions $v^*(t,x)$, $\frac{\partial v^*(t,x)}{\partial x}$, and $\frac{\partial v^*(t,x)}{\partial t}$ on Ω , respectively, as $k \rightarrow \infty$.

The uniqueness of solution to problem (5)–(8) is proved by contradiction.

Theorem 1 is proved.

From equivalence of problem (1)–(4) and (5)–(8) follows the following assertion.

Theorem 2. Suppose that the conditions i) – iv) of Theorem 1 are fulfilled.

Then nonlocal problem with integral conditions for third order system of partial differential equations (1)–(4) has a unique classical solution.

Conditions of Theorem 1 are sufficient of the well-posedness to nonlocal problem with integral solutions for third order system of partial differential equations (1)–(4).

Funding. This results are partially supported by grant of the Ministry education and science of Republic Kazakhstan No. AP 05131220 for 2018-2020 years.

REFERENCES

- [1] Ptashnyck B.I. Ill-posed boundary value problems for partial differential equations, Naukova Dumka, Kiev (1984) (in Russian).
- [2] Nakhushhev A.M. Problems with shift for a partial differential equations, Nauka, Moskow (2006). ISBN: 5-02-034076-6 (in Russian).
- [3] Kiguradze T., Lakshmikantham V. On initial-boundary value problems in bounded and unbounded domains for a class of nonlinear hyperbolic equations of the third order, Journal of Mathematical Analysis and Applications, **324**:2 (2006), 1242--1261.
DOI: 10.1016/j.jmaa.2005.12.067
- [4] Dzhokhadze O.M. The Riemann function for higher-order hyperbolic equations and systems with dominated lower terms, Differential Equations, **39**:10 (2003), 1440--1453.
DOI: 0012-2661/03/3910-1440
- [5] Andreev A.A., Yakovleva J.O. The Goursat problem for one hyperbolic system of the third order differential equations with two independent variables, Journal of Samara State Technical University, Ser. Physical and Mathematical Sciences, **15**:3(24) (2011), 35--41. DOI: <https://doi.org/10.14498/vsgtu996>
- [6] Assanova A.T., Imanchiev A.E. Solvability of multipoint-integral boundary value problem for a third-order differential equation and parametrization method, in: Functional Analysis in Interdisciplinary Applications, Springer Proceedings in Mathematics & Statistics, T.Sh. Kalmenov et al. (eds.), Springer International Publishing, Cham, **216** (2017), 113--122. https://doi.org/10.1007/978-3-319-67053-9_11
- [7] Assanova A.T., Sabalakhova A.P., Toleukhanova Z.M. On the solving of initial-boundary value problem for system of partial differential equations of the third order, News of the National Academy of Sciences of the RK. Physico-Mathematical Series. **3**:319 (2018), 67--73.
- [8] Assanova A.T., Imanchiyev A.E., Kadirbayeva Z.M. On the solvability of multipoint problem for the loaded third order partial differential equation, Mathematical Journal. **18**:1(67) (2018), 27-35.
- [9] Assanova A.T., Imanchiyev A.E. On the unique solvability of a family of multipoint-integral boundary value problems for a third order differential equation, Bulletin of the Karaganda University-Mathematics. 2(90) (2018), 25-33.
- [10] Asanova A.T., Dzhumabaev D.S. Unique solvability of the boundary value problem for systems of hyperbolic equations with data on the characteristics, Computational Mathematics and Mathematical Physics. **42**:11 (2002), 1609--1621.
- [11] Asanova A.T., Dzhumabaev D.S. Correct solvability of a nonlocal boundary value problem for systems of hyperbolic equations, Doklady Mathematics. **68**:1 (2003), 46--49.
- [12] Asanova A.T., Dzhumabaev D.S. Unique solvability of nonlocal boundary value problems for systems of hyperbolic equations, Differential Equations. **39**:10 (2003), 1414--1427. 0012-2661/03/3910-1414
- [13] Asanova A.T., Dzhumabaev D.S. Periodic solutions of systems of hyperbolic equations bounded on a plane, Ukrainian Mathematical Journal. **56**:4 (2004), 682--694.
0041-5995/04/5604-0682
- [14] Asanova A.T., Dzhumabaev D.S. Well-posed solvability of nonlocal boundary value problems for systems of hyperbolic equations, Differential Equations. **41**:3 (2005), 352--363.
0012-2661/05/4103-0352
- [15] Asanova A.T. A nonlocal boundary value problem for systems of quasilinear hyperbolic equations, Doklady Mathematics. **74**:3 (2006), 787--791.
DOI: 10.1134/S1064562406060019
- [16] Asanova A.T. On the unique solvability of a nonlocal boundary value problem with data on intersecting lines for systems of hyperbolic equations, Differential Equations. **45**:3 (2009), 385--394. DOI: 10.1134/S0012266109030082
- [17] Asanova A.T. On a boundary-value problem with data on noncharacteristic intersecting lines for systems of hyperbolic equations with mixed derivative, Journal of Mathematical Sciences (United States). **187**:4 (2012), 375--386. 1072-3374/12/1874-0375
- [18] Asanova A.T. On a nonlocal boundary-value problem for systems of impulsive hyperbolic equations, Ukrainian Mathematical Journal. **65**:3 (2013), 349--365.
0041-5995/13/6503-0349
- [19] Asanova A.T., Dzhumabaev D.S. Well-posedness of nonlocal boundary value problems with integral condition for the system of hyperbolic equations, Journal of Mathematical Analysis and Applications, **402**:1 (2013), 167--178. doi:10.1016/j.jmaa.2013.01.012
- [20] Asanova A.T. Well-posed solvability of a nonlocal boundary-value problem for systems of hyperbolic equations with impulse effects, Ukrainian Mathematical Journal. **67**:3 (2015), 333--346. DOI: 10.1007/s11253-015-1083-3

- [21] Asanova A.T. On solvability of nonlinear boundary value problems with integral condition for the system of hyperbolic equations, *Electronic Journal of Qualitative Theory of Differential Equations*. 63 (2015), 1--13. doi: 10.14232/ejqtde.2015.1.63
- [22] Asanova A.T., Imanchiev A.E. On conditions of the solvability of nonlocal multi-point boundary value problems for quasi-linear systems of hyperbolic equations, *Eurasian Mathematical Journal*. 6:4 (2015), 19--28.
- [23] Asanova A.T. Multipoint problem for a system of hyperbolic equations with mixed derivative, *Journal of Mathematical Sciences (United States)*, 212:3 (2016), 213--233.
DOI: 10.1007/s10958-015-2660-6
- [24] Asanova A.T. Criteria of solvability of nonlocal boundary-value problem for systems of hyperbolic equations with mixed derivatives, *Russian Mathematics*. 60:1 (2016), 1-17.
DOI: 10.3103/S1066369X16050017
- [25] Assanova A.T. On the solvability of nonlocal boundary value problem for the systems of impulsive hyperbolic equations with mixed derivatives, *Journal of Discontinuity, Nonlinearity and Complexity*. 5:2 (2016), 153--165. DOI: 10.5890/DNC.2016.06.005
- [26] Assanova A.T. Periodic solutions in the plane of systems of second-order hyperbolic equations, *Mathematical Notes*. 101:1 (2017), 39--47. DOI: 10.1134/S0001434617010047
- [27] Assanova A.T. Nonlocal problem with integral conditions for a system of hyperbolic equations in characteristic rectangle, *Russian Mathematics*. 61:5 (2017), 7--20.
DOI: 10.3103/S1066369X17050024
- [28] Assanova A.T. Solvability of a nonlocal problem for a hyperbolic equation with integral conditions, *Electronic Journal of Differential Equations*. 2017:170 (2017), 1-12.
- [29] Assanova A.T. An approach to solution of a nonlocal problem for a system of hyperbolic equations with integral conditions, *Nonlinear Oscillations*, 20:4 (2017), 435-450. (in Russian)
- [30] Assanova A.T., Kadirbaeva Zh. M., and Bakirova E. A. On the unique solvability of a nonlocal boundary-value problem for systems of loaded hyperbolic equations with impulsive actions, *Ukrainian Mathematical Journal*. 69:8 (2018), 1175—1195.
DOI: 10.1007/s11253-017-1424-5
- [31] Assanova A.T. On a nonlocal problem with integral conditions for the system of hyperbolic equations, *Differential Equations*. 54:2 (2018), 201–214.
DOI: 10.1134/S0012266118020076
- [32] Assanova A.T., Kadirkayeva Z.M. Periodic problem for an impulsive system of the loaded hyperbolic equations, *Electronic Journal of Differential Equations*. 2018:72 (2018), 1-8.

УДК 517.951

МРНТИ 27.31.15

А.Т. Асанова¹, Б.Ж. Алиханова², К.Ж.Назарова²

¹Институт математики и математического моделирования, Алматы, Казахстан

²Международный Казахско-Турецкий университет им. Х.А.Яссави, Туркестан, Казахстан

**КОРРЕКТНАЯ РАЗРЕШИМОСТЬ НЕЛОКАЛЬНОЙ ЗАДАЧИ
С ИНТЕГРАЛЬНЫМИ ДЛЯ СИСТЕМЫ ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ
В ЧАСТНЫХ ПРОИЗВОДНЫХ ТРЕТЬЕГО ПОРЯДКА**

Аннотация. Рассматривается нелокальная задача с интегральными условиями для системы дифференциальных уравнений в частных производных третьего порядка. Исследуются вопросы существования и единственности классического решения нелокальной задачи для системы дифференциальных уравнений в частных производных третьего порядка и предлагаются методы построения их приближенных решений. Установлены условия однозначной разрешимости нелокальной задачи для системы дифференциальных уравнений в частных производных третьего порядка. Путем введения новой неизвестной функции исследуемая задача сведена к эквивалентной задаче, состоящей из нелокальной задачи для системы гиперболических уравнений второго порядка с интегральными условиями и функциональными параметрами и интегрального соотношения. Предложены алгоритмы нахождения приближенного решения исследуемой задачи и доказана их сходимость. Получены достаточные условия существования единственного решения эквивалентной задачи с параметрами. Корректная разрешимость нелокальной задачи с интегральными условиями для системы дифференциальных уравнений в частных производных третьего порядка получены в терминах корректной разрешимости нелокальной задачи с интегральными условиями для системы гиперболических уравнений второго порядка.

Ключевые слова: дифференциальное уравнение в частных производных третьего порядка, нелокальная задача, интегральное условие, система гиперболических уравнений второго порядка, разрешимость, алгоритм.

УДК 517.951
МРНТИ 27.31.15

А.Т. Асанова¹, Б.Ж. Алиханова², К.Ж. Назарова²

¹Математика және математикалық моделдеу институты, Алматы, Қазақстан
²Қ.А.Ясауи атындағы Халықаралық қазақ-түрк университеті, Түркістан, Қазақстан

**ҮШІНШІ РЕТТІ ДЕРБЕС ТУЫНДЫЛЫ ДИФФЕРЕНЦИАЛДЫҚ
ТЕҢДЕУЛЕР ЖҮЙЕСІ ҮШІН ИНТЕГРАЛДЫҚ ШАРТТАРЫ БАР
БЕЙЛОКАЛ ЕСЕПТІҢ КОРРЕКТЛІ ШЕШІЛМДІЛІГІ**

Аннотация. Ушінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін интегралдық шарттары бар бейлокал есеп қарастырылады. Ушінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін бейлокал есептің классикалық шешімінің бар болуы мен жалғыздығы мәселелері зерттеледі және олардың жуық шешімдерін тұрғызу әдістері ұсынылады. Ушінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін интегралдық шарттары бар бейлокал есептің бірмәнді шешілімдігінің шарттары тағайындалған. Жаңа белгісіз функция енгізу арқылы зерттеліп отырған есеп гиперболалық теңдеулер жүйесі үшін интегралдық шарттары және параметрлері бар бейлокал есептен және интегралдық қатынастан тұратын пара-пар есепке келтірілген. Зерттеліп отырған есептің жуық шешімін табу алгоритмдері ұсынылған және олардың жинақтылығы дәлелденген. Параметрлері бар пара-пар есептің жалғыз шешімінің бар болуының жеткілікті шарттары алынған. Ушінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін интегралдық шарттары бар бейлокал есептің корректілі шешілімділігінің шарттары екінші ретті гиперболалық теңдеулер жүйесі үшін интегралдық шарттары бар бейлокал есептің корректілі шешілімділігі терминінде алынған.

Түйін сөздер: үшінші ретті дербес туындылы дифференциалдық теңдеу, бейлокал есеп, интегралдық шарт, екінші ретті гиперболалық теңдеулер жүйесі, шешілімділік, алгоритм.

Information about authors:

Assanova Anar Turmaganbetkyzy - the member of Editorial Board of journal "News of the NAS RK. Physico-Mathematical Series", Institute of Mathematics and Mathematical Modeling, chief scientific researcher, Doctor of Physical and Mathematical Sciences, professor, anarasanova@list.ru, assanova@math.kz, <https://orcid.org/0000-0001-8697-8920>;

Alikhanova Botakoz Zhorakhanovna - Akhmet Yassawi International Kazakh-Turkish University, teacher, bota_mktu@mail.ru, <https://orcid.org/0000-0003-2912-1300>;

Nazarova Kulgina Zharkimbekovna - Akhmet Yassawi International Kazakh-Turkish University, head teacher, gjnazarova@mail.ru, <https://orcid.org/0000-0002-2093-1879>;

МАЗМУНЫ

| | |
|---|----|
| <i>Малков Е.А., Момынов С.Б.</i> Хенон-Хейлес потенциалының фазалық бейнесі..... | 5 |
| <i>Каракеев Т.Т., Мустафаева Н.Т.</i> Ж. Бірінші түрдегі сыйыкты емес интегралды Вольтерра теңдеулерін сандық шешу әдісі..... | 10 |
| <i>Харин С. Н., Қасабек С. А., Салыбек Д., Ашимов Т.</i> Эллипсоидтік координаттардағы стефан есебі..... | 19 |
| <i>Джумабаев Д.С., Бакирова Э.А., Кадирбаева Ж.М.</i> Параметрі бар дифференциалдық теңдеулер үшін басқару есебін шешудің бір алгоритмі туралы..... | 25 |
| <i>Асанова А.Т., Алиханова Б.Ж., Назарова К.Ж.</i> Ушінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін интегралдық шарттары бар бейлоқал есептің корректілі шешілімділігі..... | 33 |
| <i>Сейтмұратов А.Ж., Tileubay С.Ш., Тоқсанова С.К., Ибрағимова Н.Ж., Досжанов Б.А., Айтимов М.Ж.</i> Қатаң шекаралармен шектелген серпімді қабат тербелісі жайлы есеп..... | 42 |
| <i>Ахметов Дж. Ш., Сейтова С.М., Тойбазаров Д.Б., Кадырбаева Г.Т., Даулеткулова А.У., Исаева Г. Б.</i> Техникалық құрылғылардың жауапкершілікті және аталған мүмкіндіктерді тиімділікті арқылы..... | 49 |
| <i>Калимолдаев М.Н., Абдилаева А.А., Ахметжанов М.А., Галиева Ф.М.</i> Электр энергетикалық жүйелерді тиімді басқару мәселесін математикалық модельдеу..... | 62 |
| <i>Харин С. Н., Қасабек С. А., Слямхан М.</i> Қөпір эрозиясының теория есебі..... | 69 |

СОДЕРЖАНИЕ

| | |
|--|----|
| <i>Малков Е.А., Момынов С.Б.</i> Фазовые портреты потенциала Хенона-Хейлеса..... | 5 |
| <i>Каракеев Т.Т., Мустафаева Н.Т.</i> Метод численного решения нелинейных интегральных уравнений Вольтерра первого рода..... | 10 |
| <i>Харин С. Н., Касабек С. А., Салыбек Д., Ашимов Т.</i> Задача стефана в эллипсоидальных координатах..... | 19 |
| <i>Джумабаев Д.С., Бакирова Э.А., Кадирбаева Ж.М.</i> Об одном алгоритме решения задачи управления для дифференциальных уравнений с параметром..... | 25 |
| <i>Асанова А.Т., Алиханова Б.Ж., Назарова К.Ж.</i> Корректная разрешимость нелокальной задачи с интегральными для системы дифференциальных уравнений в частных производных третьего порядка | 33 |
| <i>Сейтмуратов А.Ж., Тилеубай С.Ш., Токсанова С.К., Ибрагимова Н.Ж., Досжанов Б.А., Айтимов М.Ж.</i> Задача о колебании упругого слоя ограниченные жесткими границами | 42 |
| <i>Ахметов Дж.Ш., Сейтова С.М., Тойбазаров Д.Б., Кадырбаева Г.Т., Даuletкулова А.У., Исаева Г. Б.</i> Проверка технических устройств надежности через решение вероятности неисправности и неисправности..... | 49 |
| <i>Калимолдаев М.Н., Абдилаева А.А., Ахметжанов М.А., Галиева Ф.М.</i> Математическое моделирование задачи оптимального управления электроэнергетическими системами..... | 62 |
| <i>Харин С. Н., Касабек С. А., Слямхан М.</i> Задача из теории мостиковой эрозии..... | 69 |

CONTENTS

| | |
|--|----|
| <i>Malkov E.A., Momynov S.B.</i> Phase portraits of the Henon-Heiles potential..... | 5 |
| <i>Karakeev T.T., Mustafayeva N.T.</i> The method of numerical solution of nonlinear Volterra integral equations of the first kind..... | 10 |
| <i>Kharin S.N., Kassabek S.A., Salybek D., Ashymov T.</i> Stefan problem in ellipsoidal coordinates..... | 19 |
| <i>Dzhumabaev D.S., Bakirova E.A., Kadirbayeva Zh.M.</i> An algorithm for solving a control problem for a differential equation with a parameter..... | 25 |
| <i>Assanova A.T., Alikhanova B.Zh., Nazarova K.Zh.</i> Well-posedness of a nonlocal problem with integral conditions for third order system of the partial differential equations | 33 |
| <i>Seitmuratov A., Tileubay S., Toxanova S., Ibragimova N., Doszhanov B., Aitimov M.Zh.</i> The problem of the oscillation of the elastic layer bounded by rigid boundaries | 42 |
| <i>Akhmetov J.W., Seitova S.M., Toibazarov D.B., Kadyrbayeva G.T., Dauletkulova A.U., Issayeva G. B.</i> Verification of reliability technical devices through resolving probability of failure and failure..... | 49 |
| <i>Kalimoldayev M.N., Abdildayeva A.A., Akhmetzhanov M.A., Galiyeva F.M.</i> Mathematical modeling of the problem of optimal control of electric power systems..... | 62 |
| <i>Kharin S.N., Kassabek S.A., Slyamkhan M.</i> Problem from the theory of bridge erosion..... | 69 |

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www:nauka-nanrk.kz

http://www.physics-mathematics.kz

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов
Верстка на компьютере А.М. Кульгинбаевой

Подписано в печать 11.10.2018.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
4,9 п.л. Тираж 300. Заказ 5.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19