

ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

ӘЛЪ-ФАРАБИ АТЫНДАҒЫ
ҚАЗАҚ ҰЛТТЫҚ УНИВЕРСИТЕТІНІҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

КАЗАХСКИЙ НАЦИОНАЛЬНЫЙ
УНИВЕРСИТЕТ ИМЕНИ АЛЬ-ФАРАБИ

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

AL-FARABI KAZAKH
NATIONAL UNIVERSITY

ФИЗИКА-МАТЕМАТИКА СЕРИЯСЫ



СЕРИЯ ФИЗИКО-МАТЕМАТИЧЕСКАЯ



PHYSICO-MATHEMATICAL SERIES

5 (321)

ҚЫРКҮЙЕК – ҚАЗАН 2018 ж.
СЕНТЯБРЬ – ОКТЯБРЬ 2018 г.
SEPTEMBER – OCTOBER 2018

1963 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1963 ГОДА
PUBLISHED SINCE JANUARY 1963

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі **Ғ.М. Мұтанов**

Р е д а к ц и я а л қ а с ы:

Жұмаділдаев А.С. проф., академик (Қазақстан)
Кальменов Т.Ш. проф., академик (Қазақстан)
Жантаев Ж.Ш. проф., корр.-мүшесі (Қазақстан)
Өмірбаев У.У. проф. корр.-мүшесі (Қазақстан)
Жүсіпов М.А. проф. (Қазақстан)
Жұмабаев Д.С. проф. (Қазақстан)
Асанова А.Т. проф. (Қазақстан)
Бошқаев К.А. PhD докторы (Қазақстан)
Сұраған Д. корр.-мүшесі (Қазақстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Қырғыстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Белорус)
Пашаев А. проф., академик (Әзірбайжан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5543-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.
Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / physics-mathematics.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2018

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.ф.-м.н., проф. академик НАН РК **Г.М. Мутанов**

Редакционная коллегия:

Джумадильдаев А.С. проф., академик (Казахстан)
Кальменов Т.Ш. проф., академик (Казахстан)
Жантаев Ж.Ш. проф., чл.-корр. (Казахстан)
Умирбаев У.У. проф. чл.-корр. (Казахстан)
Жусупов М.А. проф. (Казахстан)
Джумабаев Д.С. проф. (Казахстан)
Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Сураган Д. чл.-корр. (Казахстан)
Quevedo Hernando проф. (Мексика),
Джунушалиев В.Д. проф. (Кыргызстан)
Вишневский И.Н. проф., академик (Украина)
Ковалев А.М. проф., академик (Украина)
Михалевич А.А. проф., академик (Беларусь)
Пашаев А. проф., академик (Азербайджан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)

«Известия НАН РК. Серия физико-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов
Министерства культуры и информации Республики Казахстан №5543-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
www.nauka-nanrk.kz / physics-mathematics.kz

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75.

E d i t o r i n c h i e f
doctor of physics and mathematics, professor, academician of NAS RK **G.M. Mutanov**

E d i t o r i a l b o a r d:

Dzhumadildayev A.S. prof., academician (Kazakhstan)
Kalmenov T.Sh. prof., academician (Kazakhstan)
Zhantayev Zh.Sh. prof., corr. member. (Kazakhstan)
Umirbayev U.U. prof. corr. member. (Kazakhstan)
Zhusupov M.A. prof. (Kazakhstan)
Dzhumabayev D.S. prof. (Kazakhstan)
Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Suragan D. corr. member. (Kazakhstan)
Quevedo Hernando prof. (Mexico),
Dzhunushaliyev V.D. prof. (Kyrgyzstan)
Vishnevskiy I.N. prof., academician (Ukraine)
Kovalev A.M. prof., academician (Ukraine)
Mikhalevich A.A. prof., academician (Belarus)
Pashayev A. prof., academician (Azerbaijan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 5543-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
www.nauka-nanrk.kz/physics-mathematics.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

<https://doi.org/10.32014/2018.2518-1726.10>

Volume 5, Number 321 (2018), 75 – 87

UDC 517.94

A. A. Shaldanbaeva¹, M.I. Akylbayev², A. Sh. Shaldanbaev³, A.Zh. Beisebaeva⁴

^{1,2}"Regional Social-Innovative University", Shymkent;

^{3,4}M.Auezov South Kazakhstan State University, Shymkent;

altima_a@mail.ru, musabek_kz@mail.ru, shaldanbaev51@mail.ru, akbope_a@mail.ru.

THE SPECTRAL DECOMPOSITION OF CAUCHY PROBLEM'S SOLUTION FOR LAPLACE EQUATION

Abstract: The spectral decomposition of Cauchy problem for Laplace equation is obtained in Krein space, and is made a regularization of a problem, using the resolvent of the corresponding operator.

Keywords: spectrum, spectral decomposition, equation with deviating argument, Hilbert-Schmidt theorem, Cauchy Problem, Laplace equation, incorrect, range.

1. Introduction.

Currently, there are different approaches to the solution of Cauchy problem for elliptic equations, which is a classic example of an ill-posed problem. All approaches basically can be divided on two large groups. One group consists of methods based on the introduction of the problem into the class of correctness by Tikhonov [1] - [3], the other are the methods using the universal regularizing algorithms, obtained by means of the parametric functional of Tikhonov [4].

It should be noted that the second group of methods received the most spread and major achievements in the practical application. In this approach, are used different variants of regularized algorithms that reduce the problem or to the solution of integral equations of the first kind, or to the representation of the desired field in the region beside or to the construction of finite-difference regularized algorithms [4] — [6].

In view of a great importance of the problem, which has applications in many fields of science and technology, and constantly rising requirements for the reliability of the results, the search for other approaches to its solution is continued. Iterative methods in recent years are becoming more widely used in the practice of solutions of various ill-posed problems of mathematical physics [7] - [9]. These methods have a number of undoubted advantages, which include simple computational schemes, their uniformity for applications with linear and nonlinear operators, the high accuracy of the solution, and so on.

An important advantage is the fact that they allow simple accounting of the essential restrictions for tasks on the solution directly in the scheme of the iterative algorithm (e.g., restrictions on non-negativity of solutions, monotonicity, and so on). In [10] was proposed a new method for solving the problem in question, based on the alternating iterative procedure, which is a consistent solution of the correct mixed boundary value problems for the original equation.

It is proved the convergence of the method and its regularizing properties. This method is general and can be extended to a wide range of similar ill-posed boundary value problems of mathematical physics. The weak point of the method is the requirement for the smoothness of the boundary, which is not always fulfilled, in particular, in our case. In this paper we propose a spectral method [11-14].

Let $\Omega = [-1, 1] \times [0, \pi]$ be a rectangle with sides

$AB: y = 0, -1 \leq x \leq 1; BC: x = 1, 0 \leq y \leq \pi; CD: y = \pi, -1 \leq x \leq 1; DA: x = -1, 0 \leq y \leq \pi$

(see fig.1)




Fig 1.

Let's consider the following Cauchy - Dirichlet problem for Poisson's equation in the region Ω :

$$Lu = u_{xx} - u_{yy} = f(x, y), \quad (1)$$

$$u|_{y=0} = 0, \quad u|_{y=\pi} = 0, \quad (2)$$

$$u|_{x=-1} = 0, \quad \frac{\partial u}{\partial x}|_{x=-1} = 0, \quad (3)$$

where $f(x, y) \in L^2(\Omega)$. This problem has been investigated previously in [11], [12] and is found that inverse operator L^{-1} exists, but is unlimited, in particular, it was shown that the "smallest" eigenvalue of the operator $A = SL$ has asymptotics

$$\lambda_{m0} = 4m^2 e^{-2m} [1 + o(1)], \quad m \rightarrow \infty,$$

where the operator S has the form $Su(x, y) = u(-x, y)$.

This work complements and refines the results of these studies.

2. Research Methods

The main idea of the method belongs to T.Sh. Kalmenov [11], and consists in the following. The operator $A = SL$ is symmetric in the space $L^2(\Omega)$, so with the original problem the boundary value problem is studied

$$SLu = Sf, \quad (1')$$

$$u|_{y=0} = 0, \quad u|_{y=\pi} = 0, \quad (2')$$

$$u|_{x=-1}, \quad \frac{\partial u}{\partial x}|_{x=-1} = 0, \quad (3')$$

where the operator S has the form, see [13] - [15].

$$Su(x, y) = u(-x, y),$$

and resembles an involution of M.G. Krein, see [16].

The following spectral problem corresponds to this boundary value problem (1') - (3')

$$Au = \lambda u,$$

$$\begin{aligned} u|_{y=0} = 0, \quad u|_{y=\pi} = 0, \\ u|_{x=-1}, \quad \frac{\partial u}{\partial x}|_{x=-1} = 0, \end{aligned}$$

where

$$A = SL,$$

or in expanded form

$$u_{xx} + u_{yy} = \lambda u(-x, y), \tag{4}$$

$$u|_{y=0} = 0, \quad u|_{y=\pi} = 0,$$

$$u|_{x=-1} = 0, \quad \frac{\partial u}{\partial x}|_{x=-1} = 0, \tag{5}$$

We solve this spectral problem (4) - (5) using the method of separation of variables, assuming

$$u(x, y) = v(x)w(y),$$

and as a result we get two spectral problems

$$a) -w''(y) = \mu w(y), \tag{6}$$

$$w(0) = 0, \quad w(\pi) = 0; \tag{7}$$

$$b) v''(x) - \mu v(x) = \lambda v(-x), \tag{8}$$

$$v(-1) = 0, \quad v'(-1) = 0. \tag{9}$$

The solution of (6) - (7) is well known and has the form $w_m(y) = \sin my$, $m = 1, 2, \dots$; an analogue of the spectral problem (8) - (9) was investigated in detail in [14], however, we will give a full and detailed study of this problem in the fourth section of the article. As a result, we have

$$Au_{mn} = \lambda_{mn}u_{mn}, \quad m = 1, 2, \dots; \quad n = 0, 1, 2, \dots$$

where $\{u_{mn}\}, m = 1, 2, \dots; \quad n = 0, 1, 2, \dots$ is complete and orthonormal system of functions in the space $L^2(\Omega)$.

Further, from the equation (1') we have

$$\bar{A}u = Sf,$$

Where \bar{A} is a closure of the operator A in the space $L^2(\Omega)$.

Hence,

$$\begin{aligned} u &= (\bar{A})^{-1}Sf = \sum_{m,n} \langle (\bar{A})^{-1}Sf, u_{mn} \rangle u_{mn} = |(\bar{A})^* = \bar{A}| = \langle Sf, (\bar{A})^{-1}u_{mn} \rangle u_{mn} \\ &= \left| Au_{mn} = \lambda_{mn}u_{mn}, \Rightarrow \bar{A}u_{mn} = \lambda_{mn}u_{mn}, \quad A \subset \bar{A}, \Rightarrow (\bar{A})^{-1}u_{mn} = \frac{u_{mn}}{\lambda_{mn}} \right| \\ &= \sum_{m,n} \frac{(Sf, u_{mn})}{\lambda_{mn}} u_{mn} = \sum_{m,n} \frac{[f, u_{mn}]}{\lambda_{mn}} u_{mn}, \end{aligned}$$

where $[f, u_{mn}] = (Sf, u_{mn})$ is the inner product of Krein's space, and $(.,.)$ is the usual inner product of the space $L^2(\Omega)$, i.e.,

$$(f, g) = \iint_{\Omega} f(x, y) \cdot \overline{g(x, y)} dx dy.$$

Therefore, we need to show a closability of operator A , an essential self-adjointness: $\bar{A} = A^*$, and reversibility: $\ker \bar{A} = \{0\}$, because all of these properties are used in derivation of the last formula

$$u(x, y) = \sum_{m,n}^{\infty} \frac{[f, u_{mn}]}{\lambda_{mn}} u_{mn}(x, y).$$

In addition, it is necessary to examine the spectrum of the operator \bar{A} .

3. Results of research.

Let $D(A)$ is a domain of definition of operator A , and $R(A)$ is a domain of its values, $\ker A$ is a kernel of the operator A , where

$$A = SL$$

$$Lu = u_{xx} + u_{yy}, \quad Su(x, y) = u(-x, y),$$

$$D(A) = \left\{ u(x, y) \in C^2(\Omega) \cap C^1(\bar{\Omega}): u|_{y=0} = 0, \quad u|_{y=\pi} = 0, u|_{x=-1}, \quad \frac{\partial u}{\partial x} \Big|_{x=-1} = 0 \right\};$$

We denote through \bar{A} the closure of the operator A in the space $L^2(\Omega)$.

The following theorem holds

Theorem 1.

- (a) A is closable, i.e. its closure exists;
- (b) A is essentially self-adjoint in the space $L^2(\Omega)$, i.e. the equality holds $(\bar{A})^* = \bar{A}$;
- (c) A is invertible, i.e. $\ker \bar{A} = \{0\}$, but the inverse operator $(\bar{A})^{-1}$ is unlimited, and has the form

$$u(x, y) = (\bar{A})^{-1}Sf(x, y) = \sum_{m,n}^{\infty} \frac{(Sf, u_{mn})}{\lambda_{mn}} u_{mn}(x, y) = \sum_{m,n}^{\infty} \frac{[f, u_{mn}]}{\lambda_{mn}} u_{mn}(x, y),$$

where $\{u_{mn}\}, m = 1, 2, \dots; n = 0, 1, 2, \dots$ are the orthonormal eigenvectors of A , and λ_{mn} are the corresponding eigenvalues;

d) $\overline{R(A)} = H = L^2(\Omega) \neq R(\bar{A})$;

i.e. the operator equation

$$\bar{A}u = Sf$$

is densely solvable in the space $L^2(\Omega)$, but not everywhere solvable. The following theorem 2 reveals the spectral properties of the operator \bar{A} .

Theorem 2. The spectrum of operator \bar{A} consists of four parts

a) The negative part:

$$-m^2 - (n\pi)^2 < \lambda_{mn}^- < -m^2 - \left(n\pi + \frac{\pi}{4}\right)^2, \quad m, n = 1, 2, \dots;$$

b) the "zero" part:

$$m^2 e^{-2\sqrt{2}m \cos u_0(m^2)} < \lambda^0(m^2) < 2(1 + \sqrt{2})m^2 e^{-2\sqrt{2}m \sin u_0(m^2)},$$

where

$$u_0(m^2) \geq u_0(1) > u_0^*(1) > 0,$$

and

$$\lim_{m \rightarrow \infty} u_0(m^2) = \frac{\pi}{4}; \quad m = 1, 2, \dots$$

c) the positive part

$$m^2 + (n\pi)^2 < \lambda_{mn}^+ < m^2 + \left(n\pi + \frac{\pi}{2}\right)^2, \quad m, n = 1, 2, \dots$$

d) the limit part, i.e. point $\lambda = 0$ belongs to the limit spectrum of the operator \bar{A} , i.e. the equality holds

$$\overline{\{\lambda_{mn}\}} \ni \{0\}$$

g) the inequalities hold $\lambda_{mn} \neq 0, m = 1, 2, \dots; n = 0, 1, 2, \dots$

Theorem 3. The boundary value problem

$$u_{xx} + u_{yy} = \lambda u(-x, y), \quad x \in (-1, 1], \quad y(x, \pi)$$

$$u|_{y=0} = 0, \quad u|_{y=\pi} = 0,$$

$$u|_{x=-1}, \quad \frac{\partial u}{\partial x}|_{x=-1} = 0,$$

has a complete and orthogonal system of eigenvectors:

$$u_{mn}(x, y) = \sin my * v_{mn}(x), \quad m = 1, 2, \dots; n = 0, 1, 2, \dots$$

$$v_{mn}(x) = K_{mn} [ch\sqrt{m^2 + \lambda_{mn}}sh\sqrt{m^2 - \lambda_{mn}}xsh\sqrt{m^2 - \lambda_{mn}}ch\sqrt{m^2 + \lambda_{mn}}x],$$

$$m = 1, 2, \dots; n = 0, 1, 2, \dots$$

where K_{mn} are the normalization coefficients, and λ_{mn} are the roots of the equation

$$th\sqrt{m^2 - \lambda}th\sqrt{m^2 + \lambda} = \frac{\sqrt{m^2 - \lambda}}{\sqrt{m^2 + \lambda}},$$

for each fixed value $m = 1, 2, \dots$

All the eigenvalues are simple, real and not equal to zero.

4. The proofs and discussion.

4.1. On the solvability.

Lemma 1. If the eigenvectors of a symmetric operator T , corresponding to non-zero eigenvalues, form an orthonormal basis of the Hilbert space H , then

- a) this operator is essentially self-adjoint;
 - b) the operator \bar{T} is reversible;
 - c) $R(\bar{T}) = H$;
 - d) $R(\bar{T}) = R(T)$,
- if and only if the inequality holds

$$|\lambda_n| \geq \varepsilon > 0, n = 1, 2, \dots$$

where $\lambda_n (n = 1, 2, \dots)$ are the eigenvalues of the operator T acting in a Hilbert space H .

4.2. On the spectrum of the operator B .

Consider in the space $L^2(-1, 1)$ the following spectral problem

$$v''(x) - \mu v(x) = \lambda v(-x), x \in (-1, 1] \tag{8}$$

$$v(-1) = 0, \quad v'(-1) = 0 \tag{9}$$

where μ is a fixed real quantity, λ is a spectral parameter.

Let $B = S\hat{L}$, where

$$\hat{L}v = v''(x) - \mu v(x), \quad Su(x) = u(-x),$$

then the spectral problem (8) - (9) takes the form

$$Bv = \lambda v; \quad v(-1) = 0, v'(-1) = 0.$$

Notice that

$$D(B) = \{v(x) \in C^2(-1, 1) \cap C^1[-1, 1]: v(-1) = 0, v'(-1) = 0\}.$$

Obviously $C_0^\infty(-1, 1) \subset D(B)$. In addition, the equality holds

$$(Bu, v) = (u, Bv), \quad \forall u, v \in D(B)$$

If $\lambda = 0$, then $v(x) = 0$ by virtue of the uniqueness of the solution of Cauchy problem, so $ker B = \{0\}$, i.e. the inverse operator B^{-1} exists, which has the following form

$$v(x) = B^{-1}f(x) = \int_{-1}^x \frac{sh\sqrt{\mu}(x-t)Sf(t)}{\sqrt{\mu}} dt,$$

for any continuous function $f(x) \in C[-1,1]$. By means of the extension theorem (see [19], C.154), we will continue this operator on the whole space $L^2(-1,1)$ as a continuous operator

$$\overline{B^{-1}}f(x) = \int_{-1}^x \frac{sh\sqrt{\mu}(x-t)Sf(t)}{\sqrt{\mu}} dt, \quad \forall f(x) \in L^2(-1,1).$$

It is obvious that the operator $\overline{B^{-1}}$ is completely continuous and self-adjoint in the space $L^2(-1,1)$. By the formula,

$$\overline{B^{-1}} = (\overline{B})^{-1}$$

we have that the operator \overline{B} is reversible. Operator $(\overline{B})^{-1}$ is completely continuous and self-adjoint in $L^2(-1,1)$.

By Hilbert-Schmidt Theorem (see [17], p. 226), for any $f(x) \in L^2(-1,1)$ the formula holds

$$\begin{aligned} (\overline{B})^{-1}f &= \sum_{n=1}^{\infty} \langle (\overline{B})^{-1}f, v_n(\mu; x) \rangle v_n(\mu; x) = \sum_{n=1}^{\infty} \langle f, (\overline{B})^{-1}v_n(\mu; x) \rangle v_n(\mu; x) \\ &= \sum_{n=1}^{\infty} \langle f, \overline{B^{-1}} v_n(\mu; x) \rangle v_n(\mu; x) = \left| \overline{B^{-1}}v_n = B^{-1}v_n = \frac{v_n(\mu; x)}{\lambda_n(\mu)} \right| \\ &= \sum_{n=1}^{\infty} \langle f, v_n(\mu; x) \rangle \frac{v_n(\mu; x)}{\lambda_n(\mu)}, \end{aligned}$$

where $\lambda_n(\mu)$ are the eigenvalues of the operator B , and $v_n(\mu; x)$ are the corresponding eigenvectors.

If $\langle f, v_n(\mu; x) \rangle = 0$, for $n = 1, 2, \dots$, then $(\overline{B})^{-1}f = 0$, hence $f = 0$, i.e. the system $\{v_n(\mu; x)\}, n = 1, 2, \dots$ is complete and orthogonal in the space $L^2(-1,1)$. We formulate the obtained results as following lemma.

Lemma 2. If $\mu = \overline{\mu}$, i.e. it is a real value, then

- the operator $(\overline{B})^{-1}$ is completely continuous and self-adjoint;
- the spectrum of \overline{B} is discrete, i.e., it has no the condensation points;
- the normalized eigenvectors of \overline{B} form the orthonormal basis of the space $L^2(-1,1)$.

Let's find the eigenfunctions of the problem (8) - (9). The general solution of equation (8) has the form

$$v(\mu, \lambda; x) = a(\mu, \lambda)sh\sqrt{\mu - \lambda}x + b(\mu, \lambda)ch\sqrt{\mu + \lambda}x, \quad (10)$$

where $a(\mu, \lambda), b(\mu, \lambda)$ are arbitrary constants.

Indeed,

$$\begin{aligned} v'(\mu, \lambda; x) &= a(\mu, \lambda)\sqrt{\mu - \lambda}ch\sqrt{\mu - \lambda}x + b(\mu, \lambda)\sqrt{\mu + \lambda}sh\sqrt{\mu + \lambda}x, \quad (11) \\ v''(\mu, \lambda; x) &= a(\mu, \lambda)(\mu - \lambda)sh\sqrt{\mu - \lambda}x + b(\mu, \lambda)(\mu + \lambda)ch\sqrt{\mu + \lambda}x \\ &= \mu\lambda(\mu, \lambda; x)[-a(\mu, \lambda)sh\sqrt{\mu - \lambda}x + b(\mu, \lambda)ch\sqrt{\mu + \lambda}x]\lambda = \mu v(\mu, \lambda; x) + \lambda v(\mu, \lambda; -x), \\ &\Rightarrow v''(\mu, \lambda; x) - \mu v(\mu, \lambda; x) = \lambda v(\mu, \lambda; -x). \end{aligned}$$

Substituting (10) - (11) into the boundary conditions (9), we have

$$\begin{cases} -ash\sqrt{\mu - \lambda} + bch\sqrt{\mu + \lambda} = 0 \\ a\sqrt{\mu - \lambda}ch\sqrt{\mu - \lambda} - b\sqrt{\mu + \lambda}sh\sqrt{\mu + \lambda} = 0 \end{cases} \quad (*)$$

This system of equations has a nontrivial solution only if its determinant $\Delta(\mu, \lambda)$ equals 0, where

$$\Delta(\mu, \lambda) = \begin{vmatrix} -sh\sqrt{\mu - \lambda} & ch\sqrt{\mu + \lambda} \\ \sqrt{\mu - \lambda}ch\sqrt{\mu - \lambda} & -\sqrt{\mu + \lambda}sh\sqrt{\mu + \lambda} \end{vmatrix}.$$

Expanding this determinant, we obtain

$$\Delta(\mu, \lambda) = \sqrt{\mu + \lambda}sh\sqrt{\mu + \lambda}sh\sqrt{\mu - \lambda} - \sqrt{\mu - \lambda}ch\sqrt{\mu - \lambda}ch\sqrt{\mu + \lambda} \tag{12}$$

If $\lambda = 0$, then

$$\Delta(\mu, 0) = \sqrt{\mu}sh^2\sqrt{\mu} - \sqrt{\mu}ch^2\sqrt{\mu} = -\sqrt{\mu}[ch^2\sqrt{\mu} - sh^2\sqrt{\mu}] = -\sqrt{\mu},$$

If in addition $\mu = 0$, then from (8) - (9) we have that $v(x) = 0$. Consequently, the value of $\lambda = 0$ is not an eigenvalue of the problem (8) - (9).

If $\lambda = \mu$, then $\Delta(\mu, \mu) = 0$, therefore, the value of $\lambda = \mu$ is probably the eigenvalue of the problem (8) - (9), to which the following eigenfunction corresponds

$$v(\mu, \mu; x) = b(\mu, \mu)ch\sqrt{2\mu}x.$$

But this function satisfies the boundary condition (9) only when the $b(\mu, \mu) = 0$, so there is no eigenvalues of the boundary value problem (8) - (9) in the segment $[-\mu, 0]$, $\mu \geq 0$.

Assuming $\mu > 0$ for definiteness, let us study the distribution of zeros of functions (12).

Lemma 3. If $\mu > \mu_0$, then the function $F(\mu, u)$ has a unique simple zero u_0 , located in the interval $0 < u_0^* < u_0 < \frac{\pi}{4}$, where

$$1 - \sqrt{2\mu_0}th\sqrt{2\mu_0} = 0, \quad \left. \frac{\partial F}{\partial u} \right|_{u=u_0^*} = 0.$$

Lemma 4. If $z = u$ a real quantity, the

a) for $0 < \mu < \mu_0$ segment $[-\mu, \mu]$ no eigenvalues;

b) when $\mu \geq \mu_0$ in interval $(0, \mu)$ will be exactly one eigenvalue $\lambda_0^+(\mu)$, which satisfies the estimate

$$\mu \cdot e^{-2\sqrt{2\mu} \cos u_0(\mu)} < \lambda_0^+(\mu) < 2\mu(1 + \sqrt{2}) \cdot e^{-2\sqrt{2\mu} \sin u_0(\mu)},$$


where

$$u_0(\mu) > u_0^*(\mu) > 0, \quad \forall \mu > \mu_0, \quad \lim_{\mu \rightarrow +\infty} u_0(\mu) = \frac{\pi}{4},$$

$$\left. \frac{\partial F}{\partial u} \right|_{u=u_0^*} = 0, \quad 1 - \sqrt{2\mu_0}th\sqrt{2\mu_0} = 0.$$

a) $0 < \mu \leq \mu_0$

$-\infty + \infty$



b) $\mu > \mu_0 > 0$

$-\infty \lambda_0^+$



fig 5.

Consequence. For all $\mu \geq 1$, the inequality holds

$$\mu \cdot e^{-2\sqrt{2\mu}} < \lambda_0^+(\mu) < 2\mu(1 + \sqrt{2})e^{-2\sqrt{2\mu} \sin u_0^*(1)}$$

where

$$\left. \frac{\partial u}{\partial u} \right|_{u=u_0^*(1)} = 0, \quad u_0^*(1) > 0.$$

Lemma 5.

a) If $0 < \mu < \mu_0$, then the eigenvalues of the spectral task

$$v''(x) - \mu v(x) = \lambda v(-x), \quad x \in (-1,1] \quad (8)$$

$$v(-1) = 0, \quad v'(-1) = 0 \quad (9)$$

consists of two series: negative

$$-\left[\mu + \left(n\pi + \frac{\pi}{2}\right)^2\right] < \lambda_n^- < -\left[\mu + \left(n\pi + \frac{\pi}{4}\right)^2\right], \quad n = 0,1,2, \dots,$$

and positive

$$\mu + (n\pi)^2 < \lambda_n^+(\mu) < \mu + \left(n\pi + \frac{\pi}{2}\right)^2, \quad n = 0,1,2 \dots$$

b) If $\mu > \mu_0$, then the third "zero" series will appear, that lays in the interval $(0, \mu)$, for which a two-sided estimate is valid

$$\mu \cdot e^{-2\sqrt{2\mu} \cos u_0(\mu)} < \lambda_0^+(\mu) < 2\mu(1 + \sqrt{2})e^{-2\sqrt{2\mu} \sin u_0(\mu)}$$

where

$$u_0(\mu) > u_0^*(\mu) > 0, \quad \forall \mu > \mu_0, \quad \lim_{\mu \rightarrow +\infty} u_0(\mu) = \frac{\pi}{4}$$

$$\left. \frac{\partial F}{\partial u} \right|_{n=u_0^*} = 0, \quad 1 - \sqrt{2\mu_0} \operatorname{th} \sqrt{2\mu_0} = 0$$

b) we will name the quantity μ_0 the threshold, it is the root of the equation

c)

$$1 - \sqrt{2\mu} \operatorname{th} \sqrt{2\mu} = 0$$

for it the assessment holds: $0,5 < \mu_0 < 0,72$, see. ([18], p.33)

From the system of equations (*) and formula (10), see. P.10 assuming

$$a_n = K_n \operatorname{ch} \sqrt{\mu + \lambda_n}, b_n = K_n \operatorname{sh} \sqrt{\mu - \lambda_n},$$

we will find the eigenfunctions of the boundary value problem (8) - (9)

$$v_n(\mu, \lambda_n, x) = K_n [\operatorname{ch} \sqrt{\mu + \lambda_n} \operatorname{sh} \sqrt{\mu - \lambda_n} x + \operatorname{sh} \sqrt{\mu - \lambda_n} \operatorname{ch} \sqrt{\mu - \lambda_n} x].$$

Lemma 6. If $\mu \geq 1$, then the spectral task

$$v''(x) - \mu v(x) = \lambda v(-x), \quad x \in (-1,1] \quad (8)$$

$$v(-1) = 0, \quad v'(-1) = 0 \quad (9)$$

has complete and orthogonal system of eigenvectors:

$$v_n(\mu, \lambda_n, x) = K_n [\operatorname{ch} \sqrt{\mu + \lambda_n} \operatorname{sh} \sqrt{\mu - \lambda_n} x + \operatorname{sh} \sqrt{\mu - \lambda_n} \operatorname{ch} \sqrt{\mu - \lambda_n} x], \quad n = 0,1,2, \dots$$

corresponding to the real eigenvalues $\lambda_n(\mu)$, $n = 0,1,2, \dots$, which are distributed as follows:

a) negative:

$$-\left[\mu + \left(n\pi + \frac{\pi}{2}\right)^2\right] < \lambda_n^-(\mu) < -\left[\mu + \left(n\pi + \frac{\pi}{4}\right)^2\right], \quad n = 0,1,2 \dots$$

b) zero:

$$\mu \cdot e^{-2\sqrt{2\mu} \cos u_0(\mu)} < \lambda_0^+(\mu) < 2\mu(1 + \sqrt{2})e^{-2\sqrt{2\mu} \sin u_0(\mu)}$$

where

$$u_0(\mu) > u_0^*(\mu) > 0, \quad \forall \mu > \mu_0, \quad \lim_{\mu \rightarrow +\infty} u_0(\mu) = \frac{\pi}{4} \text{ (monotonically)}$$

$$\frac{\partial F}{\partial u} \Big|_{u=u_0^*(\mu)} = 0, \quad 1 - \sqrt{2\mu_0} \operatorname{th} \sqrt{2\mu_0} = 0, \quad 0,5 < \mu_0 < 0,72$$

c)

positive:

$$\mu + (n\pi)^2 < \lambda_n^+(\mu) < \mu + (n\pi + \frac{\pi}{2})^2, \quad n = 1, 2, \dots$$

g) the interval $[-\mu, 0]$ remains free of eigenvalues where $\mu \geq 0$;**4.3. The proofs of theorems.**

Let's start with Theorem 3. Assuming

$$u_{mn}(x, y) = \sin my \cdot v_{mn}(x), \quad m = 1, 2, \dots$$

We divide the variables of the equation

$$u_{xx} + u_{yy} = \lambda u(-x, y), \quad x \in (-1, 1], \quad y \in (0, \pi)$$

result for $u_{mn}(x, y)$ we obtain the spectral problem

$$v''_{mn} - m^2 v_{mn}(x) = \lambda_{mn} v_{mn}(-x), \quad m = 1, 2, \dots \\ v_{mn}(-1) = 0, \quad v'_{mn}(-1) = 0.$$

By virtue of the proven Lemma 6, eigenfunctions of the spectral problem $\{v_{mn}(x)\}$, $m = 1, 2, \dots, n = 0, 1, 2, \dots$ form a complete orthogonal system in the space $L^2(-1, 1)$. Therefore, after normalization, they form an orthonormal basis of the space.

Lemma 7. If the system $\{\varphi_m(y)\}$, $m = 1, 2, \dots$ is an orthonormal basis of the space $L^2(0, \pi)$, and the system $\{\psi_{mn}(x)\}$, $m = 1, 2, \dots; n = 0, 1, 2, \dots$ for each fixed value of m is an orthonormal basis of the space $L^2(-1, 1)$, then the system

$$u_{mn}(x, y) = \psi_{mn}(x) \cdot \varphi_m(y), \quad m = 1, 2, \dots; n = 0, 1, 2, \dots$$

is an orthonormal basis of the space $L^2(\Omega)$, where $\Omega = [-1, 1] \times [0, \pi]$. See the proof [14].

In our case,

$$\varphi_m(y) = \sqrt{\frac{2}{\pi}} \sin my, \quad \psi_{mn}(x) = K_{mn} \cdot v_{mn}(x), \quad m = 1, 2, \dots; n = 0, 1, 2, \dots,$$

where K_{mn} are the normalization coefficients, hence eigenfunctions

$$u_{mn}(x, y) = K_{mn} \sin my v_{mn}(x), \quad m = 1, 2, \dots; n = 0, 1, 2, \dots$$

of the boundary value problem

$$u_{xx} + u_{yy} = \lambda u(-x, y), \quad x \in (-1, 1], \quad y \in (0, \pi) \\ u|_{y=0} = 0, \quad u|_{y=\pi} = 0, \\ u|_{x=-1}, \quad \frac{\partial u}{\partial x} \Big|_{x=-1} = 0,$$

After normalization form an orthonormal basis of the space $L^2(\Omega)$. Theorem 3 is proved. Theorem 2 is a consequence of Lemma 6, when $\mu = m^2$, $m = 1, 2, \dots$. Theorem 1 follows from Lemma 1 and Theorem 3.

5. Conclusions.

The operator \bar{A} has a spectral hatch $(-\mu_0, 0)$, where $0,5 < \mu_0 < 0,72$;

- 1) If $f(x, y) \in L^2(\Omega)$, then the solution of (1) - (3) exists if and only if
 2)

$$\sum_{m=1}^{\infty} \sum_{n=0}^{+\infty} \frac{|(Sf, u_{mn})|^2}{\lambda_{mn}^2} < +\infty,$$

where $Sf(x, y) = f(-x, y)$.

- 2) There is regularizing family of tasks, which has the form
 $\bar{A}u + \mu u = Sf$, где $0 < \mu < \mu_0$.

Proof of paragraph 3).


Let $0 < \mu < \mu_0$, we estimate the resolution of $(\bar{A} + \mu I)^{-1}$. It's obvious that
 $\bar{A}u_{mn} + \mu u_{mn} = (\lambda_{mn} + \mu)u_{mn}$, therefore $(\bar{A} + \mu I)^{-1}u_{mn} = \frac{u_{mn}}{\lambda_{mn} + \mu}$.

$$\begin{aligned} u_{\mu}(x, y) &= [\bar{A} + \mu I]^{-1}Sf \\ &= \sum_{m,n} ([\bar{A} + \mu I]^{-1}Sf, u_{mn})u_{mn} \\ &= \sum_{m,n} (Sf, [\bar{A} + \mu I]^{-1}, u_{mn})u_{mn} = \sum_{m,n} \frac{(Sf, u_{mn})}{\mu + \lambda_{mn}} u_{mn}. \end{aligned}$$

- Next, we estimate the distance from the point $-\mu$ to the spectrum of the operator \bar{A} .
 a) For the negative eigenvalues of the operator \bar{A} the following inequality holds

$$-m^2 - (n\pi)^2 < \lambda_{mn}^- < -m^2 - (n\pi - \frac{\pi}{2})^2, m, n = 1, 2, \dots,$$

Therefore
 $-\infty + \infty$



since, $0 < \mu < \mu_0 < 0,72$, then $0,72 < -\mu_0 < -\mu < 0$;


$$|\lambda_{mn}^- + \mu| > \left| -1 - \left(\frac{\pi}{2}\right)^2 + \mu \right| > \left| -1 - \left(\frac{\pi}{2}\right)^2 + \mu_0 \right| = \left| 1 + \left(\frac{\pi}{2}\right)^2 - \mu_0 \right| = 1 - \mu_0 + \left(\frac{\pi}{2}\right)^2;$$

- b) to "zero" eigenvalues $\lambda_{mn}^+ > 0$, we have

$$|-\mu - \lambda_{mn}^+| > |-\mu - 0| = |-\mu| = \mu > 0;$$

For positive eigenvalues the following inequalities hold

$$m^2 + (n\pi)^2 < \lambda_{mn}^+ < m^2 + (n\pi + \frac{\pi}{2})^2, m, n = 1, 2, \dots$$



$$|-\mu - \lambda_{mn}^+| \geq \mu + \lambda_{mn}^+ > \mu + 1 + \pi^2 > 1 + \pi^2.$$

Based on these inequalities, we estimate the resolvent of \bar{A} , at $\lambda = -\mu_0$.
From the equation,

$$(\bar{A} + \mu I)u_\mu = Sf,$$

we have

$$\begin{aligned} u_\mu(x, y) &= [\bar{A} + \mu I]^{-1} Sf \\ &= \sum_{m,n} \frac{(Sf, u_{mn})}{\lambda_{mn} + \mu} u_{mn} \\ &= \sum_{m,n} \frac{(Sf, u_{mn})}{\lambda_{mn}^- + \mu} u_{mn} + \sum_{m=1}^{\infty} \frac{(Sf, u_{m0})}{\lambda_{m0}^+ + \mu} u_{m0} = \sum_{m,n} \frac{(Sf, u_{mn})}{\lambda_{mn} + \mu} u_{mn}; \end{aligned}$$

$$\begin{aligned} \|u_\mu\|^2 &= \sum_{m,n} \frac{(Sf, u_{mn})^2}{(\lambda_{mn}^- + \mu)^2} + \sum_{m=1}^{\infty} \frac{(Sf, u_{m0})^2}{(\lambda_{m0}^+ + \mu)^2} + \sum_{m,n} \frac{(Sf, u_{mn})^2}{(\lambda_{mn}^+ + \mu)^2} \\ &\leq \frac{1}{\left[1 - \mu_0 + \left(\frac{\pi}{2}\right)^2\right]^2} \sum_{m,n} |(Sf, u_{mn})|^2 \\ &\quad + \frac{1}{\varepsilon^2} \sum_{m=1}^{\infty} |(Sf, u_{m0})|^2 + \frac{1}{1 + \pi^2} \sum_{m,n} |(Sf, u_{mn})|^2 \leq K^2(\varepsilon) \|Sf\|^2 \leq K^2(\varepsilon) \cdot \|f\|^2, \end{aligned}$$

where

$$K(\varepsilon) = \max \left\{ \frac{1}{1 - \mu_0 + \left(\frac{\pi}{2}\right)^2}, \frac{1}{\varepsilon^2}, \frac{1}{1 + \pi^2} \right\}.$$

Assume that for a given $f \in L^2(\Omega)$ there exists a solution of equation $\bar{A}u = Sf$.

then

$$\lim_{\mu \rightarrow 0} \|u_\mu - u\| = 0.$$

Indeed,

$$\begin{aligned} \|u_\mu(x, y) - u(x, y)\|^2 &= \sum_{m,n} \left(\frac{1}{\lambda_{mn} + \mu} - \frac{1}{\lambda_{mn}} \right)^2 |(Sf, u_{mn})|^2 \leq \sum_{m,n} \frac{\mu^2}{(\lambda_{mn} + \mu)^2} |(Sf, u_{mn})|^2 \\ &\leq \sum_{m,n} \frac{\mu^2}{(\lambda_{mn}^- + \mu)^2 (\lambda_{mn}^-)^2} |(Sf, u_{mn})|^2 + \sum_{m,n} \frac{\mu^2}{(\lambda_{m0}^+ + \mu)^2 (\lambda_{m0}^+)^2} |(Sf, u_{m0})|^2 \\ &\quad + \sum_{m,n}^{\infty} \frac{\mu^2}{(\lambda_{mn}^+ + \mu)^2} |(Sf, u_{mn})|^2 \\ &\leq \frac{\mu^2}{\left[1 - \mu_0 + \left(\frac{\pi}{2}\right)^2\right]^2} \sum_{m,n} \frac{|(Sf, u_{mn})|^2}{(\lambda_{mn}^-)^2} + \frac{\mu^2}{(1 + \pi^2)^2} \sum_{m,n} \frac{|(Sf, u_{mn})|^2}{(\lambda_{mn}^+)^2} \\ &\quad + \sum_{m=1}^{\infty} \frac{\mu^2}{(\lambda_{m0}^+ + \varepsilon)^2 (\lambda_{m0}^+)^2} |(Sf, u_{m0})|^2 < \end{aligned}$$

$$\begin{aligned}
 &< \mu^2 \left[\frac{1}{\left[1 - \mu_0 + \left(\frac{\pi}{2}\right)^2\right]^2} + \frac{1}{(1 + \pi^2)^2} \right] \|f\|^2 \\
 &+ \sum_{m=1}^{\infty} \frac{\mu^2 |(Sf, u_{m0})|^2}{(\lambda_{m0}^+)^2 (\lambda_{m0}^+ + \mu)} < \frac{2\mu^2}{\left[1 - \mu_0 + \left(\frac{\pi}{2}\right)^2\right]^2} + \sum_{m=1}^{\infty} \frac{\mu^2}{((\lambda_{m0}^+)^2 \lambda_{m0}^+ + \mu)^2} |(Sf, u_{m0})|^2 \\
 &\leq \left| \begin{array}{l} \lambda_{10}^+ > \lambda_{20}^+ > \dots > \lambda_{N0}^+ \\ \frac{\mu}{\lambda_{m0}^+ + \mu} < 1 \end{array} \right| \\
 &\leq \frac{2\mu^2}{\left[1 - \mu_0 + \left(\frac{\pi}{2}\right)^2\right]^2} \|f\|^2 \\
 &+ \frac{\mu^2}{(\lambda_N^+)^2} \sum_{m=1}^N |(Sf, u_{m0})|^2 \\
 &+ \sum_{N+1}^{\infty} \frac{|(Sf, u_{m0})|^2}{(\lambda_{m0}^+)^2} \\
 &< \frac{2\mu^2 \|f\|^2}{\left[1 - \mu_0 + \left(\frac{\pi}{2}\right)^2\right]^2} + \frac{\mu^2}{(\lambda_N^+)^2} \|f\|^2 \\
 &+ \sum_{N+1}^{\infty} \frac{|(Sf, u_{m0})|^2}{(\lambda_{m0}^+)^2} \leq \frac{3\mu^2}{(\lambda_N^+)^2} \|f\|^2 + \sum_{N+1}^{\infty} \frac{|(Sf, u_{m0})|^2}{(\lambda_{m0}^+)^2}.
 \end{aligned}$$

By our assumption $\sum_{m=1}^{\infty} \frac{|(Sf, u_{m0})|^2}{(\lambda_{m0}^+)^2} < +\infty$, therefore for any $\varepsilon > 0$ there exists a number $N(\varepsilon)$ such that for all $M > N(\varepsilon)$ the inequality holds

$$\sum_M^{+\infty} \frac{|(Sf, u_{m0})|^2}{(\lambda_{m0}^+)^2} < \frac{\varepsilon^2}{2}.$$

Then, at a fixed $N(\varepsilon)$, there exists a number $\delta > 0$ such that for all $0 \leq \mu < \delta$ the following inequality holds

$$\frac{3\mu^2 \|f\|^2}{(\lambda_N^+)^2} < \frac{\varepsilon^2}{2}$$

Consequently, for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\|u_\mu - u\|^2 < \varepsilon^2 \text{ i.e. } \|u_\mu - u\| < \varepsilon \text{ for all } 0 \leq \mu < \delta, \text{ as required.}$$

REFERENCES

- [1] M.M. Lavrentiev, Some ill-posed problems of mathematical physics. Novosibirsk, 1962.
- [2] L.A. Chudov, Difference methods for solutions of the Cauchy problem for the Laplace equation. Doklady Akademii Nauk SSSR. 143:4 (1962), 798_801. (in Russian)
- [3] P.N. Vabishchevich, Solution of the Cauchy problem for the Laplace equation in a doubly-connected domain. Doklady Akademii Nauk SSSR. 241:6 (1978), 1257_1260. (in Russian)
- [4] A.I. Tikhonov, V.Y. Arsenin, Methods for solving ill-posed problems. Moscow: Nauka, 1986. (in Russian)
- [5] P.N. Vabishchevich, V.V. Glasko, Yu.A. Kriksin, Solution of the Hadamard problem by means of regularizing Tikhonov algorithm. Journal of Computational Mathematics and Mathematical Physics. 19:6 (1979). 1462_1470. (in Russian)
- [6] V.N. Strakhov, S.N. Ivanov, Method of analytical continuation of the field of potential fields. Actual Problems of Computational and Applied Mathematics. Novosibirsk: Science, 1983, 195_203. (in Russian)

- [7] Methods for solving ill-posed problems and their applications. Proceedings of the All-Union Workshop. Noorus, 1982. Novosibirsk: 1982.
- [8] G.I. Vasilenko, A.M. Taratorin, Image restoration. Moscow: Radio and Communications, 1986.(in Russian)
- [9] A.V. Kryanev, Iterative method for solving ill-posed problems. Journal of Computational Mathematics and Mathematical Physics. 14:1 (1974). 25 _ 35. (in Russian)
- [10]V.A. Kozlov, V.G. Maz'ya, A.V.Fomin, An iterative method for solving the Cauchy problem for elliptic equations. Journal of Computational Mathematics and Mathematical Physics. 31:1 (1991). 64 _ 74.(in Russian)
- [11]T.Sh. Kal'menov, and U.A. Iskakova, A criterion for the strong solvability of the mixed Cauchy problem for the Laplace equation. Doklady Mathematics. 75:3 (2007). 370_373.
- [12]T.Sh. Kal'menov, and U.A. Iskakova, A method for solving the Cauchy problem for the Laplace equation. Doklady Mathematics. 78:3 (2008). 874 _ 876.
- [13]T.Sh. Kal'menov, A.Sh. Shaldanbaev, and S.T. Akhmetova, On the spectral theory of equations with deviating arguments. Mathematical Journal, - Almaty. 4:3 (2004). 41 _ 48.(in Russian)
- [14]T.Sh. Kal'menov, A.Sh. Shaldanbaev, On a criterion of solvability of the inverse problem of heat conduction. Journal of Inverse and Ill-posed Problems. 18:5, (2010). 471 _ 492.
- [15]AsylzatKopzhassarova and Abdizhakhansarsenbi, "Basis Properties of Eigenfunctions of Second-Order Differential Operators with Involution Abstract and Applied Analysis, vol. 2012, Article ID 576843, 6 pages, 2012. doi:10.1155/2012/576843
- [16]J.S. Iokhvidov, M.G. Krein, Spectral theory of operators in spaces with an indefinite metric., Proceedings of the Moscow Mathematical Society. 5, (1956). 43_72. (in Russian)
- [17]M. Reed and B. Simon, Methods of Modern Mathematical Physics. Vol.1 – Functional Analysis, Moscow, Mir, 1977. (in Russian)
- [18]Janke E., Emde F., L osch F. Special Functions. Moscow, Nauka, 1977. (in Russian).
- [19]R.D. Richtmyer, Principles of Advanced Mathematical Physics. Moscow, Mir, 1982. (in Russian)

УДК 517.94

А.А. Шалданбаева¹, М.И. Акылбаев², А.Ш. Шалданбаев³, А.Ж. Бейсебаева⁴

^{1,2}Аймақтық әлеуметтік - инновациялық университеті, Шымкент қ.
^{3,4}Оңтүстік-Қазақстан мемлекеттік университеті, Шымкент қ.

ЛАПЛАС ТЕҢДЕУІНІҢ КОШИЛІК ЕСЕБІНІҢ СПЕКТРӘЛДІК ТАРАЛЫМЫ

Аннотация. Лаплас теңдеуіне қойылған Кошидің есебінің шешімі бар екені анықталып, оның Крейннің кеңістігіндегі спектрәлді таралымы алынды. Сонан соң резольвентасы арқылы есептің жөнге келетіні көрсетілді.

Түйін сөздер: Кошидің есебі, Лапластың теңдеуі, ауытқыған аргумент, жалылық, әсіре үзіксіздік, Гилберт пен Шмидтің теоремасы.

УДК 517.94

А.А. Шалданбаева¹, М.И. Акылбаев², А.Ш. Шалданбаев³, А.Ж. Бейсебаева⁴

^{1,2}Региональный социально-инновационный университет, г. Шымкент;
^{3,4}ЮКГУ им М.О.Ауэзова, г.Шымкент.

СПЕКТРАЛЬНОЕ РАЗЛОЖЕНИЕ РЕШЕНИЯ ЗАДАЧИ КОШИ ДЛЯ УРАВНЕНИЯ ЛАПЛАСА

Аннотация. В пространстве Крейна, получено спектральное разложение решения задачи Коши уравнения Лапласа, и произведена регуляризация задачи, с помощью резольвенты соответствующего оператора.

Ключевые слова: Задача Коши, уравнение Лапласа, некорректность, спектр, отклоняющийся аргумент, самосопряженность, компактность, теорема Гильберта-Шмидта.

Information about authors:Shaldanbaeva A. A. - "Regional Social-Innovative University", Shymkent; <https://orcid.org/0000-0003-2667-3097>;Akylybayev M.I. - "Regional Social-Innovative University", Shymkent; <https://orcid.org/0000-0003-1383-4592>;Shaldanbaev A. Sh. - M.Auezov South Kazakhstan State University, Shymkent; <http://orcid.org/0000-0002-7577-8402>;Beisebaeva A.Zh. - M.Auezov South Kazakhstan State University, Shymkent; <https://orcid.org/0000-0003-4839-9156>.

МАЗМҰНЫ

<i>Малков Е.А., Момынов С.Б.</i> Хенон-Хейлес потенциалының фазалық бейнесі.....	5
<i>Каракеев Т.Т., Мустафаева Н.Т. Ж.</i> Бірінші түрдегі сызықты емес интегралды Вольтерра теңдеулерін сандық шешу әдісі.....	10
<i>Харин С. Н., Қасабек С. А., Салыбек Д., Ашимов Т.</i> Эллипсоидтік координаттардағы стефан есебі.....	19
<i>Джумабаев Д.С., Бакирова Э.А., Кадирбаева Ж.М.</i> Параметрі бар дифференциалдық теңдеулер үшін басқару есебін шешудің бір алгоритмі туралы.....	25
<i>Асанова А.Т., Алиханова Б.Ж., Назарова К.Ж.</i> Үшінші ретті дербес туындылы дифференциалдық теңдеулер жүйесі үшін интегралдық шарттары бар бейлокал есептің корректілі шешілімділігі.....	33
<i>Сейтмуратов А.Ж., Тілеубай С.Ш., Тоқсанова С.К., Ибрагимова Н.Ж., Досжанов Б.А., Айтимов М.Ж.</i> Қатаң шекаралармен шектелген серпімді қабат тербелісі жайлы есеп.....	42
<i>Ахметов Дж. Ш., Сейтова С.М., Тойбазаров Д.Б., Кадырбаева Г.Т., Даулеткулова А.У., Исаева Г. Б.</i> Техникалық құрылғылардың жауапкершілікті және аталған мүмкіндіктерді тиімділікті арқылы.....	49
<i>Калимолдаев М.Н., Абдилдаева А.А., Ахметжанов М.А., Галиева Ф.М.</i> Электр энергетикалық жүйелерді тиімді басқару мәселесін математикалық модельдеу.....	62
<i>Харин С. Н., Қасабек С. А., Слямхан М.</i> Көпір эрозиясының теория есебі.....	69
<i>Шалданбаева А.А., Ақылбаев М.И., Шалданбаев А.Ш., Бейсебаева А.Ж.</i> Лаплас теңдеуінің кошилік есебінің спектралдік таралымы.....	75

СОДЕРЖАНИЕ

<i>Малков Е.А., Момынов С.Б.</i> Фазовые портреты потенциала Хенона-Хейлеса.....	5
<i>Каракеев Т.Т., Мустафаева Н.Т.</i> Метод численного решения нелинейных интегральных уравнений Вольтерра первого рода.....	10
<i>Харин С. Н., Касабек С. А., Салыбек Д., Ашимов Т.</i> Задача Стефана в эллипсоидальных координатах.....	19
<i>Джумабаев Д.С., Бакирова Э.А., Кадирбаева Ж.М.</i> Об одном алгоритме решения задачи управления для дифференциальных уравнений с параметром.....	25
<i>Асанова А.Т., Алиханова Б.Ж., Назарова К.Ж.</i> Корректная разрешимость нелокальной задачи с интегральными для системы дифференциальных уравнений в частных производных третьего порядка	33
<i>Сейтмуратов А.Ж., Тилеубай С.Ш., Токсанова С.К., Ибрагимова Н.Ж., Досжанов Б.А., Айтимов М.Ж.</i> Задача о колебании упругого слоя ограниченные жесткими границами	42
<i>Ахметов Дж.Ш., Сейтова С.М., Тойбазаров Д.Б., Кадырбаева Г.Т., Даулеткулова А.У., Исаева Г. Б.</i> Проверка технических устройств надежности через решение вероятности неисправности и исправности.....	49
<i>Калимолдаев М.Н., Абдилдаева А.А., Ахметжанов М.А., Галиева Ф.М.</i> Математическое моделирование задачи оптимального управления электроэнергетическими системами.....	62
<i>Харин С. Н., Касабек С. А., Слямхан М.</i> Задача из теории мостиковой эрозии.....	69
<i>Шалданбаева А.А., Акылбаев М.И., Шалданбаев А.Ш., Бейсебаева А.Ж.</i> Спектральное разложение решения задачи Коши для уравнения Лапласа.....	75

CONTENTS

<i>Malkov E.A., Momynov S.B.</i> Phase portraits of the Henon-Heiles potential.....	5
<i>Karakeev T.T., Mustafayeva N.T.</i> The method of numerical solution of nonlinear Volterra integral equations of the first kind.....	10
<i>Kharin S.N., Kassabek S.A., Salybek D., Ashymov T.</i> Stefan problem in ellipsoidal coordinates.....	19
<i>Dzhumabaev D.S., Bakirova E.A., Kadirbayeva Zh.M.</i> An algorithm for solving a control problem for a differential equation with a parameter.....	25
<i>Assanova A.T., Alikhanova B.Zh., Nazarova K.Zh.</i> Well-posedness of a nonlocal problem with integral conditions for third order system of the partial differential equations.....	33
<i>Seitmuratov A., Tileubay S., Toxanova S., Ibragimova N., Doszhanov B., Aitimov M.Zh.</i> The problem of the oscillation of the elastic layer bounded by rigid boundaries.....	42
<i>Akmetov J.W., Seitova S.M., Toibazarov D.B., Kadyrbayeva G.T., Dauletkulova A.U., Issayeva G. B.</i> Verification of reliability technical devices through resolving probability of failure and failure.....	49
<i>Kalimoldayev M.N., Abdildayeva A.A., Akhmetzhanov M.A., Galiyeva F.M.</i> Mathematical modeling of the problem of optimal control of electric power systems.....	62
<i>Kharin S.N., Kassabek S.A., Slyamkhan M.</i> Problem from the theory of bridge erosion.....	69
<i>Shaldanbaeva A. A., Akylbayev M.I., Shaldanbaev A. Sh., Beisebaeva A.Zh.</i> The spectral decomposition of cauchy problem's solution for Laplace equation.....	75

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://www.physics-mathematics.kz>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы *М. С. Ахметова, Т.А. Апендиев, Д.С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 11.10.2018.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
5,7 п.л. Тираж 300. Заказ 5.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19